e at -20C	C/EBPα Antibody		Cell Signaling TECHNOLOGY®	
Store		Orders:	877-616-CELL (2355) orders@cellsignal.com	
95		Support:	877-678-TECH (8324)	
#2295		Web:	info@cellsignal.com cellsignal.com	
#		3 Trask Lane Danvers Ma	ssachusetts 01923 USA	

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IF-IC	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 42, 28	Source: Rabbit	UniProt ID: #P49715	Entrez-Gene Id: 1050			
Product Usage	Ар	plication				Dilution			
Information	We	stern Blotting				1:1000			
	Imn	nunofluorescence (I	mmunocytochemis	try)		1:50			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.							
Specificity / Sensitiv	ity C/E	C/EBPa Antibody detects endogenous levels of total C/EBPa protein.							
Source / Purificatior	sequ	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the sequence of human C/EBPα. Antibodies are purified by protein A and peptide affinity chromatography.							
Background	diffe char alter trans dept trans	CCAAT/enhancer-binding proteins (C/EBPs) are a family of transcription factors that are critical for cellular differentiation, terminal function, and inflammatory response (1). Six members of the family have been characterized (C/EBP α , β , δ , γ , ε , and ζ) and are distributed in a variety of tissues (1). Translation from alternative start codons results in two isoforms of C/EBP α (p42 and p30), which are both strong transcriptional activators (2). It has been reported that insulin and insulin-like growth factor-I stimulate the dephosphorylation of C/EBP α , which may play a key role in insulin-induced repression of GLUT4 transcription (3). Phosphorylation of C/EBP α at Thr222, Thr226, and Ser230 by GSK-3 seems to be required for adipogenesis (4).							
Background Refere	2. Lii 3. He	 Lekstrom-Hims, J. and Xanthopoulos, K.G. (1998) <i>J. Biol. Chem.</i> 273, 28545-28548. Lin, F. et al. (1993) <i>Proc. Natl. Acad. Sci. USA</i> 90, 9606-9610. Hemati, N. et al. (1997) <i>J. Biol. Chem.</i> 272, 25913-25919. Ross, S.E. et al. (1999) <i>Mol. Cell. Biol.</i> 19, 8433-8441. 							
Species Reactivity	Spec	Species reactivity is determined by testing in at least one approved application (e.g., western blot).							
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.							
Applications Key	WB:	WB: Western Blotting IF-IC: Immunofluorescence (Immunocytochemistry)							
Cross-Reactivity Ke	X : Xe	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected							
Trademarks and Patents	All ot	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.							
Limited Uses	follow condi	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.							
	appro Custo	oved, cleared, or lice omer shall not use a	ensed by the FDA on ny Product for any	r other regulatory fo diagnostic or therap	abeling statement and h preign or domestic entity peutic purpose, or other ed by CST are provided	r, for any purpose. wise in any manner			

C/EBPa Antibody (#2295) Datasheet Without Images Cell Signaling Technology

end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.