Store at -200

Ribosomal Protein L13a Antibody



Orders:

877-616-CELL (2355) orders@cellsignal.com

Support:

877-678-TECH (8324)

Web:

info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H Mk	Sensitivity: Endogenous	MW (kDa): 23	Source: Rabbit	UniProt ID: #P40429	Entrez-Gene Id: 23521	
Product Usage Information	Ар	plication		Dilution			
	We	estern Blotting			1:1000		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at $-$ 20°C. Do not aliquot the antibody.					
Specificity / Sensitiv	rity Ribo	Ribosomal Protein L13a Antibody detects endogenous levels of total ribosomal protein L13a.					
Source / Purification	-	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the sequence of human ribosomal protein L13a. Antibodies are purified by peptide affinity chromatography.					
Background	fam role trigg GAI repr	Ribosomal protein L13a (RPL13a, 60S ribosomal protein L13a) is a member of the L13 ribosomal protein family and a structural component of the 60S ribosomal subunit (1). RPL13a appears to play an important role in transcript-specific translational silencing. Interferon-y induces the phosphorylation of RPL13a and triggers the release of this protein from the 60S ribosomal subunit (2). Free RPL13a protein binds to the GAIT (interferon-y-activated inhibitor of translation) complex at the 3'-UTR of ceruloplasmin (Cp) mRNA to repress Cp expression (2). RPL13a bound to the GAIT complex interacts with eIF4G, which prevents the recruitment of 43S ribosomal subunit and results in transcript-specific translation suppression (3).					
Background Referei	2. M	 Higa, S. et al. (1999) Gene 240, 371-7. Mazumder, B. et al. (2003) Cell 115, 187-98. Kapasi, P. et al. (2007) Mol Cell 25, 113-26. 					

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with

Ribosomal Protein L13a Antibody (#2765) Datasheet Without Images Cell Signaling Technology other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection

with the Products.