

#2780 Store at -20C

PEA-15 Antibody



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R Mk	Endogenous	15	Rabbit	#Q15121	8682

Product Usage Information	Application	Dilution
	Western Blotting	1:1000
	Immunoprecipitation	1:100
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	PEA-15 Antibody detects endogenous levels of total PEA-15 protein.	
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding residue surrounding Leu60 of human PEA-15. Antibodies are purified by protein A and peptide affinity chromatography.	
Background	PEA-15 is a 15 kDa phosphoprotein expressed abundantly in astrocytes and fibroblasts as well as in tissues, including the lung and eye (1). The protein has been shown to coordinate cell growth, death, and glucose utilization (2-4). The amino-terminal DED domain of PEA-15 mediates its binding to FADD or Erk and further regulates the Erk and apoptosis signaling pathways. PEA-15 can be phosphorylated at two serine residues, Ser104 and Ser116, located within the carboxy terminus. Phosphorylation at these sites regulates binding to Erk and FADD (2,3).	
Background References	<ol style="list-style-type: none"> 1. Danziger, N. et al. (1995) <i>J. Neurochem.</i> 64, 1016-1025. 2. Krueger, J. et al. (2005) <i>Mol. Biol. Cell</i> 16, 3552-3561. 3. Renganathan, H. et al. (2005) <i>Biochem. J.</i> 390, 729-735. 4. Condorelli, G. et al. (1998) <i>EMBO J.</i> 17, 3858-3866. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting IP: Immunoprecipitation
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial</p>

purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.