

#2930 Store at -20°C

Phospho-PSD93 (Tyr340) Antibody



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP	Reactivity: R	Sensitivity: Endogenous	MW (kDa): 110	Source: Rabbit	UniProt ID: #Q15700	Entrez-Gene Id: 1740
-------------------------	------------------	----------------------------	------------------	-------------------	------------------------	-------------------------

Product Usage Information	Application Western Blotting Immunoprecipitation	Dilution 1:1000 1:50
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	Phospho-PSD93 (Tyr340) Antibody detects endogenous levels of PSD93 only when phosphorylated at Tyr340.	
Source / Purification	Antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr340 of human PSD93.	
Background	<p>Postsynaptic Density Protein 93 (PSD93) is a member of the PSD subfamily of the membrane-associated guanylate kinase (PSD-MAGUK) proteins. Structurally, it most closely resembles PSD95, consisting of an N-terminal variable segment followed by three PDZ domains, an SH3 domain, and an inactive guanylate kinase (GK) domain (1,2). PSD93 is expressed in neuronal cells and located at the synapse where it interacts with neuronal receptors and proteins including the NMDA receptor (2-4), K⁺ channels (5,6), and the AMPA receptor (7) to regulate their membrane localization and neuronal signaling. Research studies have implicated PSD93 in postsynaptic related persistent pain induction, making PSD93 a potential target for treatment of this syndrome (3).</p> <p>The phosphorylation site at Tyr340 of PSD93 was identified at Cell Signaling Technology (CST) using PhosphoScan®, CST's MS/MS platform for phosphorylation site discovery. Phosphorylation of PSD93 at Tyr340 was observed in extracts isolated from reperfusion of ischemic rat brain.</p>	
Background References	1. Brenman, J.E. et al. (1996) <i>J Neurosci</i> 16, 7407-15. 2. Kim, E. et al. (1996) <i>Neuron</i> 17, 103-13. 3. Tao, Y.X. et al. (2003) <i>J Neurosci</i> 23, 6703-12. 4. Niethammer, M. et al. (1996) <i>J Neurosci</i> 16, 2157-63. 5. Kim, E. et al. (1995) <i>Nature</i> 378, 85-8. 6. Leyland, M.L. and Dart, C. (2004) <i>J Biol Chem</i> 279, 43427-36. 7. Elias, G.M. et al. (2006) <i>Neuron</i> 52, 307-20.	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting IP: Immunoprecipitation
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.