

#2937 Store at -20C

TP/ECGF1 Antibody


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

| | | | | | | |
|----------------------------|-------------------------|-----------------------------------|----------------------------|--------------------------|-------------------------------|--------------------------------|
| Applications: WB | Reactivity: H | Sensitivity: Endogenous | MW (kDa): 50, 60 | Source: Rabbit | UniProt ID: #P19971 | Entrez-Gene Id: 1890 |
|----------------------------|-------------------------|-----------------------------------|----------------------------|--------------------------|-------------------------------|--------------------------------|

| | | |
|----------------------------------|---|---------------------------|
| Product Usage Information | Application Western Blotting | Dilution 1:1000 |
| Storage | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody. | |
| Specificity / Sensitivity | TP/ECGF1 Antibody detects endogenous levels of human TP/ECGF1 protein. | |
| Source / Purification | Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding residues surrounding Gln370 of human TP/ECGF1. Antibodies are purified by peptide affinity chromatography. | |
| Background | Thymidine phosphorylase (TP) is a platelet-derived endothelial cell growth factor (PD-ECGF) that catalyzes the formation of thymine and 2-deoxy-D-ribose-1-phosphate from thymidine and orthophosphate (1). This intracellular enzyme is capable of both promoting angiogenesis and inhibiting apoptosis. Thymidine phosphorylase catalytic activity is required for its angiogenic function (2,3). Increased expression of TP/PD-ECGF is seen in a wide variety of different solid tumors and inflammatory diseases and is often associated with poor prognosis (4,5). Alternatively, TP can activate fluorouracil derivative (DFUR) prodrugs and increase the antitumor activity of the related treatment (1,5). The use of thymidine phosphorylase as a cancer therapeutic target has been studied extensively, with emphasis on either inhibiting TP enzymatic activity or increasing enzyme induction with concomitant DFUR treatment (1,5). | |
| Background References | 1. Rooseboom, M. et al. (2004) <i>Pharmacol Rev</i> 56, 53-102. 2. Moghaddam, A. and Bicknell, R. (1992) <i>Biochemistry</i> 31, 12141-6. 3. Furukawa, T. et al. (1992) <i>Nature</i> 356, 668. 4. Toi, M. et al. (2005) <i>Lancet Oncol</i> 6, 158-66. 5. Liekens, S. et al. (2007) <i>Biochem Pharmacol</i> 74, 1555-67. | |

| | |
|-------------------------------|--|
| Species Reactivity | Species reactivity is determined by testing in at least one approved application (e.g., western blot). |
| Western Blot Buffer | IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight. |
| Applications Key | WB: Western Blotting |
| Cross-Reactivity Key | H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected |
| Trademarks and Patents | Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information. |
| Limited Uses | Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect. Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the |

end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.