

#3086 Store at -20°C

PASK (C70B2) Rabbit mAb


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R	Endogenous	170	Rabbit IgG	#Q96RG2	23178

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

PASK (C70B2) Rabbit mAb detects endogenous levels of total PASK protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to the sequence of mouse PASK.

Background

PASK is a serine/threonine kinase that contains two PAS (Per-Arnt-Sim) domains (1). Its kinase activity is up-regulated by autophosphorylation of the activation loop within its catalytic domain and is inhibited *in cis* by one of the PAS domains (1). Studies found that the yeast homolog of PASK phosphorylates and inhibits two enzymes required for glycogen biosynthesis: UDP-glucose pyrophosphorylase and glycogen synthase, resulting in the decrease of carbohydrate storage (2). Further studies showed that increased glucose levels activate PASK activity and enhance its expression in pancreatic β-cells (3). PASK is essential for the glucose-stimulated expression of preproinsulin and Pdx1, suggesting its role in the regulation of genes involved in pancreatic β-cell functions (3). PASK was shown to be critical for glucose-stimulated insulin secretion in pancreatic β-cells (4). The absence of PASK also protects animals from obesity and insulin resistance when they are fed a high-fat diet (4). These findings suggest that PASK functions as an important metabolic sensor in various cells (4).

Background References

1. Rutter, J. et al. (2001) *Proc Natl Acad Sci USA* 98, 8991-6.
2. Rutter, J. et al. (2002) *Cell* 111, 17-28.
3. da Silva Xavier, G. et al. (2004) *Proc Natl Acad Sci USA* 101, 8319-24.
4. Hao, H.X. et al. (2007) *Proc Natl Acad Sci USA* 104, 15466-71.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.