

#3224 Store at -20C

**Dvl2 (30D2) Rabbit mAb****Cell Signaling**  
TECHNOLOGY®**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R Mk	Endogenous	90 to 95	Rabbit	#O14641	1856

Product Usage Information	Application	Dilution
	Western Blotting	1:1000
	Immunoprecipitation	1:100
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	Dvl2 (30D2) Rabbit mAb detects endogenous levels of total Dvl2 protein.	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly176 of human Dvl2.	
Background	Dishevelled (Dsh) proteins are important intermediates of Wnt signaling pathways. Dsh inhibits glycogen synthase kinase-3β promoting β-catenin stabilization. Dsh proteins also participate in the planar cell polarity pathway by acting through JNK (1,2). There are three Dsh homologs, Dvl1, Dvl2 and Dvl3 in mammals. Upon treatment with Wnt proteins, Dvls become hyperphosphorylated (3) and accumulate in the nucleus (4). Dvl proteins also associate with actin fibers and cytoplasmic vesicular membranes (5) and mediate endocytosis of the Fzd receptor after Wnt protein stimulation (6). Overexpression of Dvl has been reported in certain cancers (7,8).	
Background References	<ol style="list-style-type: none"> <li>1. Logan, C.Y. and Nusse, R. (2004) <i>Annu. Rev. Cell Dev. Biol.</i> 20, 781-810.</li> <li>2. Boutros, M. and Mlodzik, M. (1999) <i>Mech. Dev.</i> 83, 27-37.</li> <li>3. Lee, J.S. et al. (1999) <i>J. Biol. Chem.</i> 274, 21464-70.</li> <li>4. Itoh, K. et al. (2005) <i>J. Biol.</i> 4, 3.1-3.12.</li> <li>5. Capelluto, D.G. et al. (2002) <i>Nature</i> 419, 726-729.</li> <li>6. Chen, W. et al. (2003) <i>Science</i> 301, 1391-1394.</li> <li>7. Okino, K. et al. (2003) <i>Oncol. Rep.</i> 10, 1219-1223.</li> <li>8. Uematsu, K. et al. (2003) <i>Oncogene</i> 22, 7218-7221.</li> </ol>	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	<b>WB:</b> Western Blotting <b>IP:</b> Immunoprecipitation
Cross-Reactivity Key	<b>H:</b> human <b>M:</b> mouse <b>R:</b> rat <b>Hm:</b> hamster <b>Mk:</b> monkey <b>Vir:</b> virus <b>Mi:</b> mink <b>C:</b> chicken <b>Dm:</b> D. melanogaster <b>X:</b> Xenopus <b>Z:</b> zebrafish <b>B:</b> bovine <b>Dg:</b> dog <b>Pg:</b> pig <b>Sc:</b> S. cerevisiae <b>Ce:</b> C. elegans <b>Hr:</b> horse <b>GP:</b> Guinea Pig <b>Rab:</b> rabbit <b>All:</b> all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit <a href="http://cellsignal.com/trademarks">cellsignal.com/trademarks</a> for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.