

#3248 Store at -20°C

PKG-1 (C8A4) Rabbit mAb


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB	H M R Mk	Endogenous	78	Rabbit IgG	#Q13976	5592

Product Usage Information	Application Western Blotting	Dilution 1:1000
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	PKG-1 (C8A4) Rabbit mAb detects endogenous levels of total PKG-1 protein, including both α and β isoforms. The antibody does not cross-react with PKG-2.	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to central residues of human PKG-1.	
Background	Cyclic GMP-dependent kinases (cGK/PKG) belong to the AGC family of serine/threonine protein kinases. In mammals, two genes encode PKG-1 and PKG-2. Alternative PKG-1 splicing yields α and β isoforms, which display tissue-specific expression patterns in humans (1). All PKG family members are activated by increased cellular cGMP, which binds to the enzyme's regulatory domain inducing a conformational change and leading to enzyme activation. cGMP levels are increased through the activation of guanylyl cyclases, a process known to occur in part through nitric oxide (NO) signaling (2). In addition to well established roles in platelet activation and smooth muscle relaxation (3), PKG signaling is important in many biological processes including cardiac contractility, axon guidance, bone growth, contraction of intestinal smooth muscle, and erectile dysfunction (4).	
Background References	1. Orstavik, S. et al. (1997) <i>Genomics</i> 42, 311-8. 2. Friebe, A. and Koesling, D. (2003) <i>Circ Res</i> 93, 96-105. 3. Lincoln, T.M. et al. (2001) <i>J Appl Physiol</i> 91, 1421-30. 4. Hofmann, F. et al. (2006) <i>Physiol Rev</i> 86, 1-23.	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the</p>

end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.