5-Lipoxygenase (C49G1) Rabbit mAb			ell Signaling ECHNOLOGY® 877-616-CELL (2355) orders@cellsignal.com	
		Support:	877-678-TECH (8324)	
#3289		Web:	info@cellsignal.com cellsignal.com	
#	3 Trask La	ne Danvers N	assachusetts 01923 USA	
For Research Use Only. Not for Use in Diagnostic Procedures.				
Applications: Reactiv WB, IP, IHC-P H	ity: Sensitivity: MW (kDa): Source/Isotype: Endogenous 78 Rabbit	UniProt ID: #P09917	Entrez-Gene Id: 240	
Product Usage	Application		Dilution	
Information	Western Blotting		1:1000	
	Immunoprecipitation		1:50	
	Immunohistochemistry (Paraffin)		1:50	
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µ 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.			
Specificity / Sensitivity	,	e (C49G1) Rabbit mAb detects endogenous levels of total 5-lipoxygenase protein. -Lipoxygenasae is very low in most tissues and cell lines, except in whole blood, bone nd macrophage cell lines.		
Species predicted to react based on 100% sequence homology:	Mouse, Rat, Monkey			
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ile168 of human 5-lipoxygenase protein.			
Background	5-Lipoxygenase (5-LO, ALOX5) is an important catalytic enzyme responsible for the biosynthesis of leukotriene LTA ₄ from arachidonic acid (1,2). Leukotriene synthesis also requires 5-lipoxygenase-activating protein (FLAP, ALOX5AP), a nuclear membrane-bound protein that binds arachidonic acid and is thought to activate 5-LO. A number of related leukotrienes (i.e. B ₄ , C ₄ , D ₄) are derived from LTA ₄ and together these lipid mediators function in immune reaction regulation. 5-LO is primarily expressed in polymorphonuclear leukocytes, peripheral blood monocytes, macrophages, and mast cells (1,3). Overexpression of 5-LO protein is seen in certain cancer cells and is associated with poor diagnosis (1,4). Depending upon the cell type, 5-LO is localized to either the cytosol or the nucleus of quiescent cells (5). Following stimulation, 5-LO translocates to the nucleus and associates with FLAP to catalyze LTA ₄ synthesis (2,3). Phosphorylation of specific residues can regulate 5-LO enzymatic activity. Phosphorylation of 5-LO at Ser523 by PKA family kinases inhibits oxygenase activity (6,7) while MAPKAP2 and ERK family kinase phosphorylation at Ser271 and Ser663 stimulates 5-LO enzymatic activity <i>in vivo</i> (8,9).			
Background References	 Woods, J.W. et al. (1995) <i>J Clin Invest</i> 95, 2035-46. Evans, J.F. et al. (2008) <i>Trends Pharmacol Sci</i> 29, 72-8. Radmark, O. et al. (2007) <i>Trends Biochem Sci</i> 32, 332-41. Chen, X. et al. (2006) <i>Curr Cancer Drug Targets</i> 6, 613-22. Werz, O. (2002) <i>Curr Drug Targets Inflamm Allergy</i> 1, 23-44. Luo, M. et al. (2004) <i>J Biol Chem</i> 279, 41512-20. Luo, M. et al. (2005) <i>J Biol Chem</i> 280, 40609-16. Werz, O. et al. (2002) <i>FASEB J</i> 16, 1441-3. Werz, O. et al. (2002) <i>J Biol Chem</i> 277, 14793-800. 			
Species Reactivity	Species reactivity is determined by testing in at least one approved	d application (e.g	., western blot).	
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.			
Applications Key	WB: Western Blotting IP: Immunoprecipitation IHC-P: Immunohistochemistry (Paraffin)			

1/1/24, 7:23 AM	5-Lipoxygenase (C49G1) Rabbit mAb (#3289) Datasheet Without Images Cell Signaling Technology
Cross-Reactivity Key	 H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.
	Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.