

#3343 Store at -20°C

## Phospho-ALK (Tyr1586) Antibody



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H	Endogenous	80 (NPM-ALK) 220 (ALK)	Rabbit	#Q9UM73	238

### Product Usage Information

#### Application

Western Blotting

#### Dilution

1:1000

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

### Specificity / Sensitivity

Phospho-ALK (Tyr1586) Antibody detects ALK only when phosphorylated at tyrosine 1586 (equivalent to Tyr646 of NPM-ALK).

### Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr1586 of human ALK. Antibodies are purified by protein A and peptide affinity chromatography.

### Background

Anaplastic lymphoma kinase (ALK) is a tyrosine kinase receptor for pleiotrophin (PTN), a growth factor involved in embryonic brain development (1-3). In ALK-expressing cells, PTN induces phosphorylation of both ALK and the downstream effectors IRS-1, Shc, PLCγ, and PI3 kinase (1). ALK was originally discovered as a nucleophosmin (NPM)-ALK fusion protein produced by a translocation (4). Investigators have found that the NPM-ALK fusion protein is a constitutively active, oncogenic tyrosine kinase associated with anaplastic lymphoma (4). Research literature suggests that activation of PLCγ by NPM-ALK may be a crucial step for its mitogenic activity and involved in the pathogenesis of anaplastic lymphomas (5).  
A distinct ALK oncogenic fusion protein involving ALK and echinoderm microtubule-associated protein like 4 (EML4) has been described in the research literature from a non-small cell lung cancer (NSCLC) cell line, with corresponding fusion transcripts present in some cases of lung adenocarcinoma. The short, amino-terminal region of the microtubule-associated protein EML4 is fused to the kinase domain of ALK (6-8).

### Background References

1. Stoica, G.E. et al. (2001) *J Biol Chem* 276, 16772-9.
2. Iwahara, T. et al. (1997) *Oncogene* 14, 439-49.
3. Morris, S.W. et al. (1997) *Oncogene* 14, 2175-88.
4. Morris, S.W. et al. (1994) *Science* 263, 1281-4.
5. Bai, R.Y. et al. (1998) *Mol Cell Biol* 18, 6951-61.
6. Rikova, K. et al. (2007) *Cell* 131, 1190-203.
7. Takeuchi, K. et al. (2008) *Clin Cancer Res* 14, 6618-24.
8. Soda, M. et al. (2007) *Nature* 448, 561-6.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

### Applications Key

**WB:** Western Blotting

### Cross-Reactivity Key

**H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.