

#3418 Store at -20°C

CTCF (D31H2) XP® Rabbit mAb



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP, IHC-P, IF-F, IF-IC, ChIP, ChIP-seq, C&R, C&T	H M R Mk	Endogenous	140	Rabbit IgG	#P49711	10664

Product Usage Information

For optimal ChIP or ChIP-seq results, use 10 µl of antibody and 10 µg of chromatin (approximately 4 x 10⁶ cells) per IP. This antibody has been validated using SimpleChIP® Enzymatic Chromatin IP Kits.

The CUT&RUN dilution was determined using CUT&RUN Assay Kit #86652.

The CUT&Tag dilution was determined using CUT&Tag Assay Kit #77552.

Application	Dilution
Western Blotting	1:1000
Immunoprecipitation	1:50
Immunohistochemistry (Paraffin)	1:640 - 1:2560
Immunofluorescence (Frozen)	1:200 - 1:800
Immunofluorescence (Immunocytochemistry)	1:400 - 1:1600
Chromatin IP	1:50
Chromatin IP-seq	1:50
CUT&RUN	1:50
CUT&Tag	1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

For a carrier free (BSA and azide free) version of this product see product #48859.

Specificity / Sensitivity

CTCF (D31H2) XP® Rabbit mAb detects endogenous levels of total CTCF protein. This antibody does not cross-react with BORIS. This antibody nonspecifically labels the lamina propria of small intestine in fixed frozen mouse tissue by immunofluorescence.

Species predicted to react based on 100% sequence homology:

Bovine, Horse

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to the carboxy terminus of the human CTCF protein.

Background

CCCTC-binding factor (CTCF) and its paralog, the Brother of the Regulator of Imprinted Sites (BORIS), are highly conserved transcription factors that regulate transcriptional activation and repression, insulator function, and imprinting control regions (ICRs) (1-4). Although they have divergent amino and carboxy termini, both proteins contain 11 conserved zinc finger domains that work in combination to bind the same DNA elements (1). CTCF is ubiquitously expressed and contributes to transcriptional regulation of cell-growth regulated genes, including *c-Myc*, *p19/ARF*, *p16/INK4A*, *BRCA1*, *p53*, *p27*, *E2F1*, and *TERT* (1). CTCF also binds to and is required for the enhancer-blocking activity of all known insulator elements and ICRs, including the H19/Igf2, Prader-Willi/Angelman syndrome, and Inactive X-Specific Transcript (XIST) anti-sense loci (5-7). CTCF DNA-binding is sensitive to DNA methylation, a mark that determines selection of the imprinted allele (maternal vs. paternal) (1). The various functions of CTCF are regulated by at least two different post-translational modifications. Poly(ADP-ribosyl)ation of CTCF is required for insulator function (8). Phosphorylation of Ser612 by protein kinase CK2 facilitates a switch of CTCF from a

transcriptional repressor to an activator at the c-Myc promoter (9). *CTCF* mutations or deletions have been found in many breast, prostate, and Wilms' tumors (10,11). Expression of BORIS is restricted to spermatocytes and is mutually exclusive of *CTCF* (3). In cells expressing BORIS, promoters of X-linked cancer-testis antigens like MAGE-A1 are demethylated and activated, but methylated and inactive in *CTCF*-expressing somatic cells (12). Like other testis specific proteins, BORIS is abnormally expressed in different cancers, such as breast cancer, and has a greater affinity than *CTCF* for DNA-binding sites, detracting from *CTCF*'s potential tumor suppressing activity (1,3,13,14).

Background References

1. Klenova, E.M. et al. (2002) *Semin Cancer Biol* 12, 399-414.
2. Klenova, E.M. et al. (1993) *Mol Cell Biol* 13, 7612-24.
3. Loukinov, D.I. et al. (2002) *Proc Natl Acad Sci USA* 99, 6806-11.
4. Mukhopadhyay, R. et al. (2004) *Genome Res* 14, 1594-602.
5. Hark, A.T. et al. (2000) *Nature* 405, 486-9.
6. Ohta, T. et al. (1999) *Am J Hum Genet* 64, 397-413.
7. Chao, W. et al. (2002) *Science* 295, 345-7.
8. Yu, W. et al. (2004) *Nat Genet* 36, 1105-10.
9. El-Kady, A. and Klenova, E. (2005) *FEBS Lett* 579, 1424-34.
10. Filippova, G.N. et al. (1998) *Genes Chromosomes Cancer* 22, 26-36.
11. Filippova, G.N. et al. (2002) *Cancer Res* 62, 48-52.
12. Vatolin, S. et al. (2005) *Cancer Res* 65, 7751-62.
13. Hong, J.A. et al. (2005) *Cancer Res* 65, 7763-74.
14. D'Arcy, V. et al. (2008) *Br J Cancer* 98, 571-9.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation **IHC-P:** Immunohistochemistry (Paraffin)
IF-F: Immunofluorescence (Frozen) **IF-IC:** Immunofluorescence (Immunocytochemistry) **ChIP:** Chromatin IP
ChIP-seq: Chromatin IP-seq **C&R:** CUT&RUN **C&T:** CUT&Tag

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
 XP is a registered trademark of Cell Signaling Technology, Inc.
 All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.