

**#3439** Store at -20°C

## Notch1 (C37C7) Rabbit mAb


**Cell Signaling**  
TECHNOLOGY®

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H	Endogenous	120, 300	Rabbit	#P46531	4851

<b>Product Usage Information</b>	<b>Application</b> Western Blotting Immunoprecipitation	<b>Dilution</b> 1:1000 1:50
<b>Storage</b>	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
<b>Specificity / Sensitivity</b>	Notch1 (C37C7) Rabbit mAb detects intracellular epitopes between 2400 and 2500 amino acids of human Notch1. It recognizes both the full-length (~300 kDa) and the NTM region (~120 kDa), which consists of a short extracellular juxtamembrane peptide, a transmembrane sequence and the intracellular domain (NICD). The antibody cannot detect the extracellular (ligand-binding) domain of Notch1 following cleavage at the S2 site by ADAM-type metalloproteases.	
<b>Source / Purification</b>	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gln2501 of human Notch1.	
<b>Background</b>	Notch proteins (Notch1-4) are a family of transmembrane receptors that play important roles in development and the determination of cell fate (1). Mature Notch receptors are processed and assembled as heterodimeric proteins, with each dimer comprised of a large extracellular ligand-binding domain, a single-pass transmembrane domain, and a smaller cytoplasmic subunit (Notch intracellular domain, NICD) (2). Binding of Notch receptors to ligands of the Delta-Serrate-Lag2 (DSL) family triggers heterodimer dissociation, exposing the receptors to proteolytic cleavages; these result in release of the NICD, which translocates to the nucleus and activates transcription of downstream target genes (3,4).	
<b>Background References</b>	1. Artavanis-Tsakonas, S. et al. (1999) <i>Science</i> 284, 770-6. 2. Chan, Y.M. and Jan, Y.N. (1998) <i>Cell</i> 94, 423-6. 3. Schroeter, E.H. et al. (1998) <i>Nature</i> 393, 382-6. 4. Rand, M.D. et al. (2000) <i>Mol Cell Biol</i> 20, 1825-35.	

<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
<b>Western Blot Buffer</b>	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
<b>Applications Key</b>	<b>WB:</b> Western Blotting <b>IP:</b> Immunoprecipitation
<b>Cross-Reactivity Key</b>	<b>H:</b> human <b>M:</b> mouse <b>R:</b> rat <b>Hm:</b> hamster <b>Mk:</b> monkey <b>Vir:</b> virus <b>Mi:</b> mink <b>C:</b> chicken <b>Dm:</b> D. melanogaster <b>X:</b> Xenopus <b>Z:</b> zebrafish <b>B:</b> bovine <b>Dg:</b> dog <b>Pg:</b> pig <b>Sc:</b> S. cerevisiae <b>Ce:</b> C. elegans <b>Hr:</b> horse <b>GP:</b> Guinea Pig <b>Rab:</b> rabbit <b>All:</b> all species expected
<b>Trademarks and Patents</b>	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
<b>Limited Uses</b>	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.