Pyk2 (5E2) Mouse mAb



Orders:

877-616-CELL (2355) orders@cellsignal.com

Support:

877-678-TECH (8324)

Web:

info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures. Dogotivity

Applications: I WB, IP	Reactivity: H M	Sensitivity: Endogenous	MW (kDa): 116	Source/Isotype: Mouse IgG2a	UniProt ID: #Q14289	Entrez-Gene Id: 2185	
Product Usage Information	Ар	plication			Dilution		
	We	estern Blotting			1:1000		
	Imr	munoprecipitation			1:50		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.					
Specificity / Sensitiv	,	Pyk2 (5E2) Mouse mAb detects endogenous levels of total Pyk2 protein. It does not cross-react with other related proteins.					
Source / Purification	-	Monoclonal antibody is produced by immunizing animals with a GST-Pyk2 fusion protein containing residues surrounding the carboxy-terminal of human Pyk2.					
Background	strue from the	Protein tyrosine kinase Pyk2, also called CAKβ, RAFTK and CADTK, is a nonreceptor tyrosine kinase structurally related to focal adhesion kinase (FAK) (1-4). Pyk2 is predominantly expressed in cells derived from hematopoietic lineages and in the central nervous system. Pyk2 is one of the signaling mediators for the G-protein-coupled receptors and MAP kinase signaling pathway. It plays an important role in cell spreading and migration (5-7).					
Background Referer	2. Lu 3. S 4. Yı 5. D 6. W	 Avraham, S. et al. (1995) J. Biol. Chem. 270, 27742-27751. Lev, S. et al. (1995) Nature 376, 737-745. Sasaki, H. et al. (1995) J. Biol. Chem. 270, 21206-21219. Yu, H. et al. (1996) J. Biol. Chem. 271, 29993-29998. Duong, L.T. et al. (2001) J Biol Chem 276, 7484-92. Watson, J.M. et al. (2001) J Biol Chem 276, 3536-42. Tang, H. et al. (2002) J Biol Chem 277, 5441-7. 					

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the Pyk2 (5E2) Mouse mAb (#3480) Datasheet Without Images Cell Signaling Technology end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection

with the Products.