

#3630 Store at -20C

RalBP1 (I33) Antibody



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H M R Mk	Endogenous	95	Rabbit	#Q15311	10928

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

RalBP1 (I33) Antibody detects endogenous levels of total RalBP1 protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ile33 of human RalBP1 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

The RalA binding protein 1 (RalBP1 or RLIP76) was originally identified as a GTP-RalA associated protein that acted as a downstream RalA effector in regulating Ral-Ras signaling (1). RalBP1 interacts with RalA and the endocytosis protein REPS2 (POB1) through its carboxy-terminal Ral binding domain. RalBP1 has an intrinsic GTPase activating function and interacts with Cdc42 through its centrally located Rho-GAP domain (1-3). A protein complex containing RalBP1/POB1/RalA regulates endocytosis of membrane receptors (4). RalBP1 also functions as a non-ABC transporter that catalyzes the ATP-dependent transport of numerous xenobiotics, including glutathione conjugates and some chemotherapeutic agents. RalBP1 transporter activity may play an important role in detoxification, drug resistance and the stress response (5-7). Increased expression of RalBP1 protein is associated with some forms of cancer and regression of cancer xenografts results from RalBP1 inhibition (8,9). Evidence to date suggests that RalBP1 may be a promising therapeutic target for cancer therapy.

Background References

1. Cantor, S.B. et al. (1995) *Mol Cell Biol* 15, 4578-84.
2. Matsubara, K. et al. (1997) *FEBS Lett* 410, 169-74.
3. Ikeda, M. et al. (1998) *J Biol Chem* 273, 814-21.
4. Nakashima, S. et al. (1999) *EMBO J* 18, 3629-42.
5. Awasthi, Y.C. et al. (2007) *Curr Drug Metab* 8, 315-23.
6. Awasthi, S. et al. (2002) *Drug Metab Dispos* 30, 1300-10.
7. Awasthi, S. et al. (2008) *Clin Cancer Res* 14, 4372-7.
8. Hudson, M.E. et al. (2007) *Proc Natl Acad Sci USA* 104, 17494-9.
9. Singhal, S.S. et al. (2007) *Cancer Res* 67, 4382-9.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.