Store at -20C

MST3 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: Reactivity: Sensitivity: MW (kDa): Source: **UniProt ID:** Entrez-Gene Id: WB HMRMk Endogenous 50 Rabbit #Q9Y6E0 8428 **Product Usage Application** Dilution Information Western Blotting 1:1000 Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at -**Storage** 20°C. Do not aliquot the antibody.

Specificity / Sensitivity MST3 Antibody recognizes endogenous levels of total MST3 protein.

Source / Purification Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asp336 of human MST3 protein. Antibodies are purified by protein A and peptide

affinity chromatography.

Mammalian sterile-20-like (MST) kinases are upstream regulators of mitogen-activated protein kinase **Background**

(MAPK) signaling pathways that regulate multiple biological processes, including apoptosis, morphogenesis, cell migration, and cytoskeletal rearrangements (1). This group of serine/threonine kinases includes a pair of closely related proteins (MST1, MST2) that are functionally distinct from the more distantly related MST3 and MST4 kinases. All four MST kinases share a conserved amino-terminal kinase domain and carboxy-terminal regulatory and interaction domains (1-3). At least three of these kinases (MST1-3) promote apoptosis and are activated by caspase cleavage followed by nuclear translocation of the active kinase. MST1/2 kinases play a key role in the Hippo signaling pathway, an evolutionarily conserved program that controls organ size by regulating cell proliferation, apoptosis, and

stem cell self renewal (4).

Mammalian Sterile 20-like kinase 3 (MST3, STK24) is ubiquitously expressed as a longer MST3b isoform and a shorter MST3a protein lacking a portion of the amino-terminal region (5). The widely expressed MST3a protein regulates apoptosis and cell motility, as well as neuronal migration during CNS development (6.7). MST3 phosphorylates and activates the NDR protein kinases that regulate cell cycle progression and cell morphology (8). Autophosphorylation of MST3 at Thr178 is required for in vitro kinase activity, and alteration of this residue inhibits MST3 regulation of cell migration in vivo (7). The brainspecific MST3b protein is activated by nerve growth factor or inosine and localizes to neurons where it

helps regulate axon growth and regeneration (9).

1. Dan, I. et al. (2001) Trends Cell Biol 11, 220-30. **Background References**

2. Creasy, C.L. et al. (1996) J Biol Chem 271, 21049-53.

3. Lee, K.K. and Yonehara, S. (2002) J Biol Chem 277, 12351-8.

4. Zhao, B. et al. (2011) Nat Cell Biol 13, 877-83.

5. Zhou, T.H. et al. (2000) J Biol Chem 275, 2513-9.

6. Tang, J. et al. (2014) J Neurosci 34, 7425-36.

7. Lu, T.J. et al. (2006) J Biol Chem 281, 38405-17.

8. Stegert, M.R. et al. (2005) Mol Cell Biol 25, 11019-29.

9. Lorber, B. et al. (2009) Nat Neurosci 12, 1407-14.

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS,

0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

Cross-Reactivity Key

WB: Western Blotting

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Limited Uses

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.