34 Store at -20C

Phospho-PLD1 (Ser561) Antibody



Orders:

877-616-CELL (2355) orders@cellsignal.com

Support:

877-678-TECH (8324)

Web:

info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 120	Source: Rabbit	UniProt ID: #Q13393	Entrez-Gene Id: 5337	
Product Usage Information	Ар	plication			Dilution		
	We	estern Blotting			1:1000		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity / Sensitivity Phospho-PLD1 (Ser561) Antibody detects endogenous levels of PLD1 only when phosphorylated at s 561. This antibody does not cross-react with phosphorylated PLD2.					sphorylated at serine		
Source / Purificat	to re	Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser561 of human PLD1. Antibodies are purified by protein A and peptide affinity chromatography.					
Background	chol Two sma	Phosphatidylcholine-specific phospholipase D (PLD) hydrolyzes phosphatidylcholine (PC) to produce choline and phosphatidic acid (PA). PA is the precursor of the second messenger, diacylglycerol (DAG). Two isoforms of PLD (PLD1 and PLD2) have been identified so far. Both are regulated by protein kinases, small GTPases and Ca2+ (1). PLD1 is phosphorylated at Ser2, Ser561, and Thr147 by PKC (2,3). Phosphorylation at Thr147 and Ser561 regulates PLD1 activity (3).					
Background Refe	2. K	 Exton, J.H. (1999) Biochim. Biophys. Acta. 1439, 121-133. Kim, Y. et al. (1999) Biochemistry 38, 10344-51. Kim, Y. et al. (2000) J. Biol. Chem. 275, 13621-13627. 					

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse

Phospho-PLD1 (Ser561) Antibody (#3834) Datasheet Without Images Cell Signaling Technology

engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.