e at -20C	ADAM9 (D64B5) Rabbit mAb				
Store a		Orders:	877-616-CELL (2355) orders@cellsignal.com		
-51		Support:	877-678-TECH (8324)		
415		Web:	info@cellsignal.com cellsignal.com		
#		3 Trask Lane Danvers	Massachusetts 01923 USA		

Applications: WB	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 100-115, 75-80	Source/Isotype: Rabbit IgG	UniProt ID: #Q13443	Entrez-Gene Id: 8754	
Product Usage Information	•	plication estern Blotting			Dilution 1:1000		
Storage	Sup	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliguot the antibody.					
Specificity / Sens		ADAM9 (D64B5) Rabbit mAb detects endogenous levels of total ADAM9 protein.					
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Val611 of human ADAM9 protein.					
Background	sign influ dom disir The at le with distr show has alter	The ADAM (A Disintegrin and A Metalloprotease) family of multidomain membrane proteins influences cell signaling and adhesion by shedding cell surface proteins such as cytokines and growth factors, by influencing cell adhesion to the extracellular matrix (ECM), and by directly remodeling the ECM. Conserved domains in ADAM family members include a prodomain, a zinc-dependent metalloprotease domain, a disintegrin domain, a cysteine-rich domain, an EGF-like sequence, and a short cytoplasmic tail (1,2). The prodomain is thought to aid in protein folding. Disintegrin and cysteine-rich domains mediate adhesion, at least in part, through binding to integrins. Phosphorylation of the cytoplasmic tail as well as its interaction with other signaling proteins may influence intra- and extracellular signaling (1). ADAM9 is widely distributed and has been shown to affect migration in skin keratinocytes (3,4). Research studies have shown that ADAM9 is overexpressed in prostate cancer (5), pancreatic cancer (6), gastric cancer (7), and has been linked to invasion and metastasis in small cell lung cancer (8). Research has also shown that an alternatively spliced short (50 kDa) form of ADAM9 containing protease activity is involved in tumor cell invasion (9).					
Background Refe	2. S 3. F 4. Z 5. F 6. G 7. C 8. S	•	Blobel, C.P. (1999) (2002) EMBO J 21 (7) J Biol Chem 28 (2008) Eur Urol 54 (2004) Br J Cance al. (2005) Int J Onc (04) Cancer Res 64	2, 30785-93. 4, 1097-106. 57 90, 1053-8. 50/ 26, 17-24. 4, 4190-6.			
Species Reactivit	t y Spec	ies reactivity is det	ermined by testing	in at least one approv	ed application (e.g., w	estern blot).	
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.					
Applications Key	WB:	WB: Western Blotting					
Cross-Reactivity	X : Xe	 H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected 					
Trademarks and Patents	All ot	Signaling Technology is a trademark of Cell Signaling Technology, Inc. ther trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more mation.					
Limited Uses							
the lunu colleignel e	om/datachaat is	n2productId=41	E1Cimages-00	protocol-0			

ADAM9 (D64B5) Rabbit mAb (#4151) Datasheet Without Images Cell Signaling Technology

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.