USP28 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H M	Sensitivity: Transfected Only	MW (kDa): 135	Source: Rabbit	UniProt ID: #Q96RU2	Entrez-Gene Id: 57646
Product Usage	Ар	Application		Dilution		
Information	We	estern Blotting			1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity / Sens	sitivity USF	USP28 Antibody detects transfected levels of USP28 protein.				
Source / Purifica		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala11 of human USP28. Antibodies are purified by peptide affinity chromatography.				
Background	deu OTU bind inclu thro	Ubiquitinating enzymes (UBEs) catalyze protein ubiquitination, a reversible process countered by deubiquitinating enzymes (DUB) action (1,2). There are five DUB subfamilies including the USP, UCH, OTU, MJD and JAMM enzymes. USP28 is an important member of the USP subfamily (3). USP28 can bind to and deubiquinate several target proteins in DNA-damage pathway, resulting in their stability, including p53BP1 and Chk2 (4). USP28 also plays an important role in Myc related signaling by binding through FBW7 α to Myc . It catalyzes the deubiquitinylation of Myc, thereby promoting its stabilization and contributing to tumor-cell growth (5).				
Background Ref	2. N 3. V 4. Z	 Nijman, S.M. et al. (2005) Cell 123, 773-86. Nalepa, G. et al. (2006) Nat Rev Drug Discov 5, 596-613. Valero, R. et al. (2001) Genome Biol 2, RESEARCH0043. Zhang, D. et al. (2006) Cell 126, 529-42. Popov, N. et al. (2007) Nat Cell Biol 9, 765-74. 				

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial

USP28 Antibody (#4217) Datasheet Without Images Cell Signaling Technology

purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.