

#4481 Store at -20°C

Phospho-PTPα (Tyr789) Antibody


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R Mk	Endogenous	145	Rabbit	#P18433	5786

Product Usage Information	Application Western Blotting Immunoprecipitation	Dilution 1:1000 1:50
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	Phospho-PTPα (Tyr789) Antibody detects endogenous levels of PTPα only when phosphorylated at Tyr789. This antibody does not cross-react with other phosphorylated receptor tyrosine phosphatases.	
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr789 of human PTPα. Antibodies are purified by protein A and peptide affinity chromatography.	
Background	PTPα (PTPRA) is a transmembrane receptor tyrosine phosphatase implicated in the regulation of Src family kinases during the G2 to mitosis entry point. Two identified splice variants differ in the size of the extracellular region; the shorter form appears to be ubiquitously expressed while the larger protein is more limited in distribution (1). The cytoplasmic region of PTPα contains two putative catalytic domains. One phosphatase domain (D1) exhibits catalytic activity while the other (D2) may regulate phosphatase activity by allowing receptor dimer formation (2,3). PTPα is a physiological regulator of Src and Src family kinases (4). Constitutive phosphorylation of the carboxy-terminal Tyr789 of PTPα is essential for dephosphorylation of Src at Tyr527. Phosphorylation of PTPα at this residue also allows binding of the Grb2 inhibitor, restricting PTPα activation of Src (5,6). PKC-mediated phosphorylation of the PTP at Ser180 and Ser204 also increases PTPα phosphatase activity (7).	
Background References	1. Kapp, K. et al. (2007) <i>Genes Cells</i> 12, 63-73. 2. Blanchetot, C. et al. (2002) <i>J Biol Chem</i> 277, 47263-9. 3. Krueger, N.X. et al. (1990) <i>EMBO J</i> 9, 3241-52. 4. den Hertog, J. et al. (1993) <i>EMBO J</i> 12, 3789-98. 5. Zheng, X.M. et al. (2000) <i>EMBO J</i> 19, 964-78. 6. Zheng, X.M. et al. (2002) <i>J Biol Chem</i> 277, 21922-9. 7. Tracy, S. et al. (1995) <i>J Biol Chem</i> 270, 10587-94.	
Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).	
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.	
Applications Key	WB: Western Blotting IP: Immunoprecipitation	
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected	
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.	
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in	

writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.