

#4640 Store at -20°C

KSR1 Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H M R Mk	Endogenous	100	Rabbit	#Q8IVT5	8844

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

KSR1 Antibody detects endogenous levels of total KSR1 protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human KSR1 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

KSR1 (kinase suppressor of Ras) was identified from a genetic screen in *Drosophila* and *C. elegans* as a component of the Ras signaling pathway (1). KSR1 has a putative carboxy-terminal kinase domain that lacks a key Lys residue for phospho-group transfer. Although reports indicate that ceramide and EGF activate KSR1 (2,3), other evidence demonstrates that KSR1 regulates Raf in a kinase-independent manner (4,5). It is now widely accepted that KSR1 functions as a scaffold that binds MEK1/2 and 14-3-3 protein constitutively and binds ERK1/2 in a Ras activation-dependent manner (1,5,6). HSP70/HSP90 and p50 Cdc37 associate with the KSR1 complex to ensure its stability (5). Multiple phosphorylation sites have been identified: Ser297 and Ser392 mediate 14-3-3 binding, and putative MAPK phosphorylation sites include Thr260, Thr274 and Ser443 (6). C-TAK1 (Cdc25C-associated kinase 1) binds and phosphorylates KSR1 at Ser392 in quiescent cells (7). In response to stimuli, Ser392 is dephosphorylated by PP2A, which leads to ERK1/2 association and allows the KSR1 complex to translocate from cytosol to membrane, where the MAPK pathway is activated (8). IMP, a Ras-responsive E3 ubiquitin ligase, is also involved in interaction with KSR1 and may regulate its localization and stability (9). Very high expression levels of KSR1 inhibit MAPK signaling, whereas physiological levels promote MAPK signaling, indicating that the scaffold protein can turn signaling "on" or "off" depending on the scaffold concentration (10).

Background References

1. Morrison, D.K. (2001) *J. Cell Sci.* 114, 1609-1612.
2. Zhang, Y. et al. (1997) *Cell* 89, 63-72.
3. Xing, H.R. and Kolesnick, R. (2001) *J. Biol. Chem.* 276, 9733-9741.
4. Michaud, N. R. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94, 12792-12796.
5. Stewart, S. et al. (1999) *Mol. Cell. Biol.* 19, 5523-5534.
6. Muller, J. et al. (2001) *Mol. Cell* 8, 983-993.
7. Cacace, A. M. et al. (1999) *Mol. Cell. Biol.* 19, 229-240.
8. Ory, S. et al. (2003) *Curr. Biol.* 13, 1356-1364.
9. Matheny, S. A. et al. (2004) *Nature* 427, 256-260.
10. Kortum, R.L. and Lewis, R.E. (2004) *Mol Cell Biol* 24, 4407-16.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.