| e at -20C | DHX29 (G63) Antibody | | Cell Signaling | |
|------------|----------------------|----------------------|--|--|
| Store at | | Orders: | 877-616-CELL (2355) orders@cellsignal.com | |
| t 8 | | Support: | 877-678-TECH (8324) | |
| #4648 | | Web: | info@cellsignal.com cellsignal.com | |
| # | | 3 Trask Lane Danvers | Massachusetts 01923 USA | |

For Research Use Only. Not for Use in Diagnostic Procedures.

| Applications: WB | Reactivity: H M R Mk | Sensitivity: Endogenous | MW (kDa): 155 | Source: Rabbit | UniProt ID: #Q7Z478 | Entrez-Gene Id: 54505 | | |
|------------------------------|--|---|--|--|--|--|--|--|
| Product Usage Information | Ар | plication | | | Dilution | | | |
| mormation | We | estern Blotting | | | 1:1000 | | | |
| Storage | | Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody. | | | | | | |
| Specificity / Sensit | t ivity DH> | DHX29 (G63) Antibody detects endogenous levels of total DHX29 protein. | | | | | | |
| Source / Purificatio | sequ | Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to a sequence around Gly63 of human DHX29 protein. Antibodies are purified by protein A and peptide affinity chromatography. | | | | | | |
| Background | subf func mRI UTF | DHX29 is an ATP-dependent RNA helicase that belongs to the DEAD-box helicase family (DEAH subfamily). DHX29 contains one central helicase and one helicase at the carboxy-terminal domain (1). Its function has not been fully established but DHX29 was recently shown to facilitate translation initiation on mRNAs with structured 5' untranslated regions (2). DHX29 binds 40S subunits and hydrolyzes ATP, GTP, UTP, and CTP. Hydrolysis of nucleotide triphosphates by DHX29 is strongly stimulated by 43S complexes and is required for DHX29 activity in promoting 48S complex formation (2). | | | | | | |
| Background Refer | | e la Cruz, J. et al. (1 isareva, V.P. et al. (| , | | | | | |
| Species Reactivity | Spec | ies reactivity is dete | ermined by testing i | n at least one appro | oved application (e.g., we | estern blot). | | |
| Western Blot Buffe | | IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight. | | | | | | |
| Applications Key | WB: | Western Blotting | | | | | | |
| Cross-Reactivity K | X : Xe | H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected | | | | | | |
| Trademarks and Patents | All ot | | ng Technology is a trademark of Cell Signaling Technology, Inc. demarks are the property of their respective owners. Visit cellsignal.com/trademarks for more | | | | | |
| Limited Uses | follov cond | Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect. | | | | | | |
| | appro Custr that o end-u thera purpo other | oved, cleared, or lic omer shall not use a conflicts with its labe user and solely for r peutic purposes, or ose, requires a sepa wise transfer or ma | ensed by the FDA of any Product for any eling statement. Pro- esearch and develor any purchase of P arate license from O ke available any Pro- | or other regulatory for diagnostic or thera oducts sold or licens opment uses. Any us roduct for resale (al- CST. Customer shall oduct to any third pa | abeling statement and har preign or domestic entity peutic purpose, or othen sed by CST are provided se of Product for diagno one or as a component) (a) not sell, license, loa arty, whether alone or in ercial products, (b) not c | r, for any purpose. wise in any manner for Customer as the stic, prophylactic or or other commercial n, donate or combination with | | |

DHX29 (G63) Antibody (#4648) Datasheet Without Images Cell Signaling Technology

engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.