

#4728 Store at -20C

β2-Chimerin (2E3) Rat mAb



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB	H M R	Endogenous	47	Rat IgG2a	#P52757	1124

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

β2-Chimerin (2E3) Rat mAb recognizes endogenous levels of total β2-chimerin protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with full-length recombinant human β2-Chimerin.

Background

Chimerins are a family of GTPase-activating proteins (GAPs) that facilitate GTP hydrolysis by the small GTPase Rac, rendering it inactive and regulating cell shape, spreading and motility. Regulation of chimerin proteins occurs in response to growth factor receptor or G-protein coupled receptor activation followed by phospholipase C activation. Chimerins are among the growing number of phorbol ester and diacylglycerol (DAG) effector molecules that do not belong to the PKC family of isoenzymes (reviewed in 1,2). β2-chimerin is highly expressed in brain and pancreas, and its expression is down-regulated in malignant gliomas (3). β2-chimerin is also down-regulated in breast cancer, and its expression causes GAP activity-dependent cell cycle arrest in MCF-7 breast cancer cells (4). Signaling from the epidermal growth factor receptor (EGFR) activates β2-chimerin and allows its association with Rac1 at the plasma membrane (5). Also in response to EGF, diacylglycerol kinase (DGK) γ interacts with β2-chimerin, promotes its translocation to the plasma membrane, and regulate its GAP activity (6).

Background References

1. Yang, C. and Kazanietz, M.G. (2007) *Biochem J.* 403, 1-12.
2. Brose, N. and Rosenmund, C. (2002) *J. Cell Sci.* 115, 4399-4411.
3. Yuan, S. et al. (1995) *Cancer Res.* 55, 3456-3461.
4. Yang, C. et al. (2005) *J. Biol. Chem.* 280, 24363-24370.
5. Wang, H. et al. (2006) *EMBO J.* 25, 2062-2074.
6. Yasuda, S. et al. (2007) *FEBS Lett.* 581, 551-557.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse **GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.