

**#4869** Store at -20°C

# HSP60 (D85) Antibody


**Cell Signaling**  
TECHNOLOGY®

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IF-IC, FC-FP	H M R Mk Dm	Endogenous	60	Rabbit	#P10809	3329

Product Usage Information	Application	Dilution
	Western Blotting	1:1000
	Immunofluorescence (Immunocytochemistry)	1:50
	Flow Cytometry (Fixed/Permeabilized)	1:50
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	HSP60 (D85) Antibody detects endogenous levels of total HSP60 protein. This antibody does not cross-react with other HSPs.	
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide surrounding Asp85 of human HSP60. Antibodies are purified by protein A and peptide affinity chromatography.	
Background	In both prokaryotic and eukaryotic cells the misfolding and aggregation of proteins during biogenesis and under conditions of cellular stress are prevented by molecular chaperones (1-3). HSP60 has primarily been known as a mitochondrial protein that is important for folding key proteins after import into the mitochondria (4). Research studies have shown that a significant amount of HSP60 is also present in the cytosol of many cells, and that it is induced by stress, inflammatory and immune responses, and autoantibodies correlated with Alzheimer's, coronary artery diseases, MS, and diabetes (5-8).	
Background References	<ol style="list-style-type: none"> <li>1. Hartl, F.U. (1996) <i>Nature</i> 381, 571-579.</li> <li>2. Bukau, B. and Horwich, A.L. (1998) <i>Cell</i> 92, 351-366.</li> <li>3. Hartl, F.U. and Hayer-Hartl, M. (2002) <i>Science</i> 295, 1852-1858.</li> <li>4. Jindal, S. et al. (1989) <i>Mol. Cell Biol.</i> 9, 2279-2283.</li> <li>5. Itoh, H. et al. (2002) <i>Eur. J. Biochem.</i> 269, 5931-5938.</li> <li>6. Gupta, S. and Knowlton, A.A. <i>J. Cell Mol. Med.</i> 9, 51-58.</li> <li>7. Deocaris, C.C. et al. (2006) <i>Cell Stress Chaperones</i> 11, 116-128.</li> <li>8. Lai, H.C. et al. (2007) <i>Am. J. Physiol. Endocrinol. Metab.</i> 292, E292-E297.</li> </ol>	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	<b>WB:</b> Western Blotting <b>IF-IC:</b> Immunofluorescence (Immunocytochemistry) <b>FC-FP:</b> Flow Cytometry (Fixed/Permeabilized)
Cross-Reactivity Key	<b>H:</b> human <b>M:</b> mouse <b>R:</b> rat <b>Hm:</b> hamster <b>Mk:</b> monkey <b>Vir:</b> virus <b>Mi:</b> mink <b>C:</b> chicken <b>Dm:</b> D. melanogaster <b>X:</b> Xenopus <b>Z:</b> zebrafish <b>B:</b> bovine <b>Dg:</b> dog <b>Pg:</b> pig <b>Sc:</b> S. cerevisiae <b>Ce:</b> C. elegans <b>Hr:</b> horse <b>GP:</b> Guinea Pig <b>Rab:</b> rabbit <b>All:</b> all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. Alexa Fluor is a registered trademark of Life Technologies Corporation. All other trademarks are the property of their respective owners. Visit <a href="http://cellsignal.com/trademarks">cellsignal.com/trademarks</a> for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.