

#5142 Store at -20°C

Loading Control Antibody Sampler Kit


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

1 Kit (5 x 20 microliters)

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
COX IV (3E11) Rabbit mAb	4850	20 µl	17 kDa	Rabbit IgG
β-Tubulin (9F3) Rabbit mAb	2128	20 µl	55 kDa	Rabbit IgG
Histone H3 (D1H2) XP® Rabbit mAb	4499	20 µl	17 kDa	Rabbit IgG
GAPDH (D16H11) XP® Rabbit mAb	5174	20 µl	37 kDa	Rabbit IgG
β-Actin (D6A8) Rabbit mAb	8457	20 µl	45 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

The Loading Control Antibody Sampler Kit contains antibodies to a variety of housekeeping proteins. The kit contains enough primary and secondary antibodies to perform two western blots per primary antibody.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Background

Housekeeping proteins perform numerous basic functions within the cell and are constitutively expressed at high levels in a variety of tissues and cell types. Western blot analysis commonly uses housekeeping proteins such as β-actin, COX IV, GAPDH, histone H3 and the α- and β-tubulins as loading controls. Actin is a ubiquitous protein and a major component of the eukaryotic cytoskeleton. Actin exists mainly as the F-actin fibrous polymer (1). Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the phosphorylation of glyceraldehyde-3-phosphate during glycolysis. Recent work has demonstrated that GAPDH plays roles in apoptosis (2), gene expression (3), and nuclear transport (4). Globular tubulin subunits made up of α- and β-tubulin heterodimers are the building blocks of microtubules, one of three types of cytosolic fibers that comprise the cytoskeleton (5). Histone proteins, including histone H3, make up the primary building block of chromatin known as nucleosomes. Modulation of the chromatin structure plays an important role in the regulation of transcription in eukaryotes (6). Cytochrome c oxidase (COX) is a hetero-oligomeric enzyme consisting of 13 subunits localized to the inner mitochondrial membrane (7-9). It is the terminal enzyme complex in the respiratory chain, catalyzing the reduction of protons across the mitochondrial inner membrane to drive ATP synthesis (10).

Background References

1. Condeelis, J. (2001) *Trends Cell Biol* 11, 288-93.
2. Hara, M.R. and Snyder, S.H. *Cell Mol Neurobiol* (2006) 26, 527-38.
3. Zheng, L. et al. (2003) *Cell* 114, 255-66.
4. Bae, B.I. et al. (2006) *Proc Natl Acad Sci USA* 103, 3405-9.
5. Westermann, S. and Weber, K. (2003) *Nat Rev Mol Cell Biol* 4, 938-47.
6. Workman, J.L. and Kingston, R.E. (1998) *Annu Rev Biochem* 67, 545-79.
7. Ostermeier, C. et al. (1996) *Curr Opin Struct Biol* 6, 460-6.
8. Capaldi, R.A. et al. (1983) *Biochim Biophys Acta* 726, 135-48.
9. Kadenbach, B. et al. (2000) *Free Radic Biol Med* 29, 211-21.
10. Barrientos, A. et al. (2002) *Gene* 286, 53-63.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.