Plexin A2 (D42B5) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cell signal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 212	Source/Isotype: Rabbit IgG	UniProt ID: #O75051	Entrez-Gene ld: 5362
Product Usage Information	Αŗ	oplication		Dilution		
	We	estern Blotting		1:1000		
	lm	munoprecipitation		1:50		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20° C. Do not aliquot the antibody.				
Specificity / Sensitivity		Plexin A2 (D42B5) Rabbit mAb detects endogenous levels of total Plexin A2 protein. This antibody also cross-reacts with unidentified proteins at various molecular weights.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly185 of human Plexin A2.				
Background	suc con anc fun trig anc Ser mig	Class 3 secreted semaphorin (Sema3A) is a chemorepellent that acts upon a wide variety of axons. As such, it induces a dramatic redistribution and depolymerization of actin filaments that results in growth cone collapse. Plexins are single membrane-spanning signaling proteins encompassing Plexin A1, A2, A3, and A4. Plexins form a complex with neuropilin-1 and -2 and the cell adhesion protein L1 to form a functional semaphorin receptor (1,2). The GTPase Rnd1 binds to the cytoplasmic domain of Plexin A1 to trigger cytoskeletal collapse. In contrast, the GTPase RhoD blocks Rnd1-mediated Plexin A1 activation and repulsion of sympathetic axons by Sema3A (3). Sema6A is also a ligand for Plexin A2. Both Sema6A and Plexin A2 knock-out mice have a granule cell migration defect, where cells remain in the molecular layer. Furthermore, Plexin A2 also controls nucleus—centrosome coupling which modulates cell migration (4).				
1. Pasterkamp, R.J. and Kolodkin, A.L. (2003) Curr Opin Neurobiol 13, 79-89. 2. Takahashi, T. and Strittmatter, S.M. (2001) Neuron 29, 429-39. 3. Zanata, S.M. et al. (2002) J Neurosci 22, 471-7. 4. Renaud, J. et al. (2008) Nat Neurosci 11, 440-9.						

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

Species Reactivity

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Plexin A2 (D42B5) Rabbit mAb (#5658) Datasheet Without Images Cell Signaling Technology

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.