

#5727 Store at -20°C

MEKK3 (D36G5) Rabbit mAb



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R Mk	Endogenous	78	Rabbit IgG	#Q99759	4215

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

MEKK3 (D36G5) Rabbit mAb recognizes endogenous levels of total MEKK3 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro310 of human MEKK3 protein.

Background

MAP kinase kinase kinase (MEKK3 or MAP3K3) is a serine/threonine protein kinase that activates SAPK and ERK via phosphorylation and activation of their respective MAP kinase kinases, SEK and MEK1/2 (1,2). MEKK3 also stimulates MEK5 via activation of ERK5/BMK1, which is at least partly regulated by a direct interaction between MEK5 and MEKK3 via p67phox-Bem1p (PB1) protein-protein interaction domains found in both proteins (3,4). MEKK3 modulates NF-κB activation in response to a variety of agonists including TNFα, LPS, IL-1 and LPA (5-9). Despite reports showing that phosphorylation of MEKK3 at Ser526 within the activation loop is necessary for kinase activation (10-12), at least one study suggests that dual phosphorylation at Thr516 and Ser520 is required for LPA-stimulated IKKβ/NF-κB activation (13). Phosphorylation at Thr294 appears to negatively regulate MEKK3 by promoting 14-3-3β binding and inhibition of the kinase activity (12). Phosphorylation of MEKK3 at Thr294 is diminished upon treatment of cells with LPS or TNFα, further suggesting an inhibitory role for this site (12).

Background References

- Blank, J.L. et al. (1996) *J Biol Chem* 271, 5361-8.
- Ellinger-Ziegelbauer, H. et al. (1997) *J Biol Chem* 272, 2668-74.
- Nakamura, K. and Johnson, G.L. (2003) *J Biol Chem* 278, 36989-92.
- Nakamura, K. et al. (2010) *J Biol Chem* 285, 2077-89.
- Yang, J. et al. (2001) *Nat Immunol* 2, 620-4.
- Zhao, Q. and Lee, F.S. (1999) *J Biol Chem* 274, 8355-8.
- Samanta, A.K. et al. (2004) *J Biol Chem* 279, 7576-83.
- Huang, Q. et al. (2004) *Nat Immunol* 5, 98-103.
- Sun, W. et al. (2009) *Cell Signal* 21, 1488-94.
- Zhang, D. et al. (2006) *EMBO J* 25, 97-107.
- Fritz, A. et al. (2006) *J Biol Chem* 281, 6236-45.
- Matitau, A.E. and Scheid, M.P. (2008) *J Biol Chem* 283, 13261-8.
- Sun, W. et al. (2010) *J Biol Chem* 285, 7911-18.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.