

#5887 Store at -20°C

γ Secretase Antibody Sampler Kit


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

1 Kit (4 x 20 microliters)

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
Nicastrin (D38F9) Rabbit mAb	5665	20 µl	110, 120 kDa	Rabbit IgG
PEN2 (D6G8) Rabbit mAb	8598	20 µl	13 kDa	Rabbit IgG
Presenilin 1 (D39D1) Rabbit mAb	5643	20 µl	22 (CTF) kDa	Rabbit IgG
Presenilin 2 (D30G3) Rabbit mAb	9979	20 µl	23 (CTF) kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

The γ Secretase Antibody Sampler Kit provides an economical means of evaluating components of the gamma secretase complex. The kit contains enough primary and secondary antibodies to perform two western miniblots experiments.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Background

The γ secretase protease complex interacts with and cleaves intramembrane substrates as an essential function for regulation of intracellular signaling and cell-cell interactions. This multiprotein complex is comprised of four integral membrane proteins, Presenilin, Nicastrin, Aph-1, and PEN2, all of which are essential for complete proteolytic activity (1). Presenilin 1 and presenilin 2 are transmembrane proteins belonging to the presenilin family. Mutation of presenilin genes has been linked to early onset of Alzheimer disease, probably due to presenilin's associated γ-secretase activity for amyloid-β protein processing (2,3). Endogenous presenilin mainly exists in a heterodimeric complex formed from the endoproteolytically processed amino-terminal (34 kDa) and carboxy-terminal (~20, 22, 23 kDa) fragments (CTF) (3,4). Nicastrin is a transmembrane glycoprotein serving as an essential component of the γ-secretase complex (5,6). Nicastrin protein is physically associated with presenilin and plays an important role in stabilization and correct localization of presenilin to the membrane-bound γ-secretase complex (7). Nicastrin also serves as a docking site for γ-secretase substrates such as APP and Notch, directly binding to them and presenting them properly to γ-secretase to ensure the correct cleavage process (6,8). Presenilin Enhancer 2 (PEN2) is a small integral membrane glycoprotein that contains two recognized transmembrane domains. Both the N- and C-terminal domains are oriented into the lumen of the endoplasmic reticulum (9). PEN2 is an important part of the γ-secretase complex as its knock down results in reduced amounts of the complex, resulting in a loss of γ-secretase activity (10).

Background References

- Hansson, C.A. et al. (2004) *J Biol Chem* 279, 51654-60.
- Haass, C. and De Strooper, B. (1999) *Science* 286, 916-9.
- Kimberly, W.T. et al. (2000) *J Biol Chem* 275, 3173-8.
- Kim, T.W. et al. (1997) *J Biol Chem* 272, 11006-10.
- Yu, G. et al. (2000) *Nature* 407, 48-54.
- Esler, W.P. et al. (2002) *Proc Natl Acad Sci U S A* 99, 2720-5.
- Kopan, R. and Goate, A. (2002) *Neuron* 33, 321-4.
- Chen, F. et al. (2001) *Nat Cell Biol* 3, 751-4.
- Sala Frigerio, C. et al. (2005) *J Neurol* 252, 1033-6.
- Steiner, H. et al. (2002) *J Biol Chem* 277, 39062-5.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.