Conjugate)

Store at +4C

85 85

**Cell Signaling** Phospho-Syk (Tyr525/526) (C87C1) Rabbit mAb (PE TECHNOLOGY® Orders: 877-616-CELL (2355) orders@cellsignal.com Support: 877-678-TECH (8324)

> Web: info@cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

cellsignal.com

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: Reactiv FC-FP H	ity: Sensitivity: Source/Isotype: Endogenous Rabbit IgG	UniProt ID:Entrez-Gene Id:#P434056850
Product Usage Information	Application	Dilution
	Flow Cytometry (Fixed/Permeabilized)	1:50
Storage	Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 n antibodies. Protect from light. Do not freeze.	ng/ml BSA. Store at 4°C. Do not aliquot the
Specificity / Sensitivity	Phospho-Syk (Tyr525/526) (C87C1) Rabbit mAb (PE Conjugate only when phosphorylated at Tyr525/526 of human Syk or Tyr5 protein when singly phosphorylated at Tyr526 of human Syk or react with other tyrosine-phosphorylated protein tyrosine kinase	19/520 of mouse Syk. It also detects Syk Tyr520 of mouse Syk. It does not cross-
Species predicted to react based on 100% sequence homology:	Mouse, Rat	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a residues surrounding Tyr525/526 of human Syk protein.	synthetic phosphopeptide corresponding to
Product Description	This Cell Signaling Technology antibody is conjugated to phyco flow cytometry analysis in human cells. The antibody is expecte reactivity as the unconjugated Phospho-Syk (Tyr525/526) (C87	d to exhibit the same species cross-
Background	Syk is a protein tyrosine kinase that plays an important role in intracellular signal transduction in hematopoietic cells (1-3). Syk interacts with immunoreceptor tyrosine-based activation motifs (ITAMs) located in the cytoplasmic domains of immune receptors (4). It couples the activated immunoreceptors to downstream signaling events that mediate diverse cellular responses, including proliferation, differentiation, and phagocytosis (4). There is also evidence of a role for Syk in nonimmune cells and investigators have indicated that Syk is a potential tumor suppressor in human breast carcinomas (5). Tyr323 is a negative regulatory phosphorylation site within the SH2-kinase linker region in Syk. Phosphorylation at Tyr323 provides a direct binding site for the TKB domain of Cbl (6,7). Tyr352 of Syk is involved in the association of PLCy1 (8). Tyr525 and Tyr526 are located in the activation loop of the Syk kinase domain; phosphorylation at Tyr525/526 of human Syk (equivalent to Tyr519/520 of mouse Syk) is essential for Syk function (9).	
Background References	<ol> <li>Cheng, A.M. and Chan, A.C. (1997) <i>Curr Opin Immunol</i> 9, 52</li> <li>Kurosaki, T. (1997) <i>Curr Opin Immunol</i> 9, 309-18.</li> <li>Chu, D.H. et al. (1998) <i>Immunol Rev</i> 165, 167-80.</li> <li>Turner, M. et al. (2000) <i>Immunol Today</i> 21, 148-54.</li> <li>Coopman, P.J. et al. (2000) <i>Nature</i> 406, 742-7.</li> <li>Deckert, M. et al. (1998) <i>J Biol Chem</i> 273, 8867-74.</li> <li>Rao, N. et al. (2001) <i>EMBO J</i> 20, 7085-95.</li> <li>Law, C.L. et al. (1996) <i>Mol Cell Biol</i> 16, 1305-15.</li> <li>Zhang, J. et al. (2000) <i>J Biol Chem</i> 275, 35442-7.</li> </ol>	8-33.
Species Reactivity	Species reactivity is determined by testing in at least one approv	red application (e.g., western blot).
Applications Key	FC-FP: Flow Cytometry (Fixed/Permeabilized)	

1/1/24, 12:50 PM Phospho-Syk (Tyr525/526) (C87C1) Rabbit mAb (PE Conjugate) (#6485) Datasheet Without Images Cell Si	
Cross-Reactivity	<ul> <li>Key</li> <li>H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster</li> <li>X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse</li> <li>GP: Guinea Pig Rab: rabbit All: all species expected</li> </ul>
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.
	Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.