

#7649 Store at -20C

Phospho-Stat1 (Tyr701) (D4A7) Rabbit mAb



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP, IF-IC, FC-FP, ChIP, ChIP-seq	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 84, 91	Source/Isotype: Rabbit IgG
---	-----------------------------	-----------------------------------	----------------------------	--------------------------------------

Product Usage Information

For optimal ChIP and ChIP-seq results, use 5 µl of antibody and 10 µg of chromatin (approximately 4 x 10⁶ cells) per IP.

This antibody has been validated using SimpleChIP® Enzymatic Chromatin IP Kits.

Application	Dilution
Western Blotting	1:1000
Immunoprecipitation	1:50
Immunofluorescence (Immunocytochemistry)	1:50
Flow Cytometry (Fixed/Permeabilized)	1:50 - 1:200
Chromatin IP	1:100
Chromatin IP-seq	1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

For a carrier free (BSA and azide free) version of this product see product #88211.

Phospho-Stat1 (Tyr701) (D4A7) Rabbit mAb recognizes endogenous levels of Stat1 protein only when phosphorylated at Tyr701.

Species predicted to react based on 100% sequence homology:

Monkey

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Tyr701 of human Stat1 protein.

Background

The Stat1 transcription factor is activated in response to a large number of ligands (1) and is essential for responsiveness to IFN-α and IFN-γ (2,3). Phosphorylation of Stat1 at Tyr701 induces Stat1 dimerization, nuclear translocation, and DNA binding (4). Stat1 protein exists as a pair of isoforms, Stat1α (91 kDa) and the splice variant Stat1β (84 kDa). In most cells, both isoforms are activated by IFN-α, but only Stat1α is activated by IFN-γ. The inappropriate activation of Stat1 occurs in many tumors (5). In addition to tyrosine phosphorylation, Stat1 is also phosphorylated at Ser727 through a p38 mitogen-activated protein kinase (MAPK)-dependent pathway in response to IFN-α and other cellular stresses (6). Serine phosphorylation may be required for the maximal induction of Stat1-mediated gene activation.

Background References

1. Heim, M.H. (1999) *J Recept Signal Transduct Res* 19, 75-120.
2. Durbin, J.E. et al. (1996) *Cell* 84, 443-50.
3. Meraz, M.A. et al. (1996) *Cell* 84, 431-42.
4. Ihle, J.N. et al. (1994) *Trends Biochem Sci* 19, 222-7.
5. Frank, D.A. (1999) *Mol Med* 5, 432-56.
6. Wen, Z. et al. (1995) *Cell* 82, 241-50.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation **IF-IC:** Immunofluorescence (Immunocytochemistry)
FC-FP: Flow Cytometry (Fixed/Permeabilized) **ChIP:** Chromatin IP **ChIP-seq:** Chromatin IP-seq

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
Alexa Fluor is a registered trademark of Life Technologies Corporation.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.