

#8192 Store at -20C

DDX3 (D19B4) Rabbit mAb


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB	H M R Mk	Endogenous	75	Rabbit IgG	#O00571	1654

Product Usage Information	Application Western Blotting	Dilution 1:1000
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	DDX3 (D19B4) Rabbit mAb recognizes endogenous levels of total DDX3 protein.	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human DDX3 protein.	
Background	The DEAD box family of RNA helicases is characterized in part by a common D-E-A-D amino acid motif. The family is composed of a growing number of proteins found in a wide range of organisms from bacteria to mammals. DEAD helicases have distinct biological functions in RNA metabolism and ribonucleoprotein (RNP) processing (reviewed in 1,2). DDX3 is a DEAD box family RNA helicase with diverse cellular functions. DDX3 is required for nuclear export of HIV-1 viral transcripts, possibly in a complex with the viral Rev protein and host cofactor CRM1 (3). DDX3 is required for hepatitis C virus (HCV) RNA replication (4) and its expression is downregulated in hepatitis B virus (HBV) associated hepatocellular carcinoma (HCC) (5). Recent evidence suggests that DDX3 functions as a tumor suppressor protein. Its expression inhibits tumor cell colony formation and increases expression of the cdk inhibitor p21 Waf1/Cip1. Low DDX3 expression has been shown in HCC (5,6), and aberrant subcellular localization occurs in many squamous cell carcinomas (6). Reduced DDX3 expression in cultured cells causes a diminished dependence on serum for cell proliferation and changes in cyclin D1 and p21 Waf1/Cip1 expression (5). DDX3 is phosphorylated at Thr204 and Thr323 by the mitotic cyclin dependent kinase, cyclin B/cdc2. This phosphorylation is thought to cause a loss of DDX3 function and a concomitant repression of ribosome biogenesis and translation in mitosis (7).	
Background References	1. Rocak, S. and Linder, P. (2004) <i>Nat Rev Mol Cell Biol</i> 5, 232-41. 2. Linder, P. (2006) <i>Nucleic Acids Res</i> 34, 4168-80. 3. Yedavalli, V.S. et al. (2004) <i>Cell</i> 119, 381-92. 4. Ariumi, Y. et al. (2007) <i>J Virol</i> 81, 13922-6. 5. Chang, P.C. et al. (2006) <i>Oncogene</i> 25, 1991-2003. 6. Chao, C.H. et al. (2006) <i>Cancer Res</i> 66, 6579-88. 7. Sekiguchi, T. et al. (2007) <i>Biochem Biophys Res Commun</i> 356, 668-73.	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses**DDX3 (D19B4) Rabbit mAb (#8192) Datasheet Without Images Cell Signaling Technology**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.