Store at -200

DDX6/RCK (D26C11) Rabbit mAb



Orders:

877-616-CELL (2355) orders@cellsignal.com

Support:

877-678-TECH (8324)

Web:

info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Applications: WB	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 54	Source/Isotype: Rabbit IgG	UniProt ID: #P26196	Entrez-Gene Id 1656
Product Usage	Application		Dilution			
Information	We	estern Blotting		1:1000		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and le 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Specificity / Sensitivity		DDX6/RCK (D26C11) Rabbit mAb recognizes endogenous levels of total DDX6/RCK protein.				
Species predicte react based on 1 sequence homol	.00%	opus				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala130 of human DDX6/RCK protein.				
Background		DDX6, also known as RCK and p54, was identified as a proto-oncogene product and is a member of the ATP-dependent DEAD box helicase family (1,2). This protein interacts with translation initiation factor eIF4E in the cytoplasmic P-bodies (3) and represses mRNA translation (4). DDX6 is a component of the miRNA induced silencing complex (miRISC) and interacts with Argonaute 1 (Ago1) and Argonaute 2 (Ago2) proteins <i>in vitro</i> and <i>in vivo</i> (5), functioning in miRNA-mediated translational repression (5).				

Background References

- 1. Akao, Y. et al. (1995) Cancer Res 55, 3444-9.
- 2. Weston, A. and Sommerville, J. (2006) Nucleic Acids Res 34, 3082-94.
- 3. Andrei, M.A. et al. (2005) RNA 11, 717-27.
- 4. Minshall, N. and Standart, N. (2004) Nucleic Acids Res 32, 1325-34.
- 5. Chu, C.Y. and Rana, T.M. (2006) PLoS Biol 4, e210.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Depletion of DDX6 leads to the disruption of cytoplasmic P-bodies indicating that it is required for P-body

Applications Key

WB: Western Blotting

formation (5).

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and **Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

DDX6/RCK (D26C11) Rabbit mAb (#8988) Datasheet Without Images Cell Signaling Technology

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.