#9099 store at -200

α-Tubulin (11H10) Rabbit mAb (HRP Conjugate)



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H M R Mk Dm Z B Pg	Sensitivity: Endogenous	MW (kDa): 52	Source/Isotype: Rabbit IgG	UniProt ID: #P68363	Entrez-Gene Id 10376	
Product Usage	App	Application			Dilution		
Information	Wes	Western Blotting			1:1000		
Storage				12 mM sodium phosphate (pH 7.4) dibasic, 2 mg/ml BSA, and iquot the antibodies.			
Specificity / Sensitivity		α -Tubulin (11H10) Rabbit mAb (HRP Conjugate) detects endogenous levels of total α -tubulin protein and does not cross-react with recombinant β -tubulin protein.					
Species predicted react based on 10 sequence homolo	0%						
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to the amino terminus of human α -tubulin protein.					
Product Description	perox	This Cell Signaling Technology antibody is conjugated to the carbohydrate groups of horseradish peroxidase (HRP) via its amine groups. The HRP conjugated antibody is expected to exhibit the same species cross-reactivity as the unconjugated antibody (α-Tubulin (11H10) Rabbit mAb #2125).					
MW (kDa)		52					
Background	and ir tubuli nucle media meml move	The cytoskeleton consists of three types of cytosolic fibers: microtubules, microfilaments (actin filaments), and intermediate filaments. Globular tubulin subunits comprise the microtubule building block, with α/β -tubulin heterodimers forming the tubulin subunit common to all eukaryotic cells. γ -tubulin is required to nucleate polymerization of tubulin subunits to form microtubule polymers. Many cell movements are mediated by microtubule action, including the beating of cilia and flagella, cytoplasmic transport of membrane vesicles, chromosome alignment during meiosis/mitosis, and nerve-cell axon migration. These movements result from competitive microtubule polymerization and depolymerization or through the actions of microtubule motor proteins (1).					
Background Refe	rences 1. We	1. Westermann, S. and Weber, K. (2003) Nat Rev Mol Cell Biol 4, 938-47.					
Species Reactivity	y Specie	es reactivity is dete	ermined by testing	g in at least one approve	ed application (e.g., we	estern blot).	
Western Blot Buff		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.					
Applications Key	WB:	Western Blotting					

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more

X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

GP: Guinea Pig Rab: rabbit All: all species expected

information.

Cross-Reactivity Key

Trademarks and

Patents

1/1/24. 11:53 AM

Limited Uses

α-Tubulin (11H10) Rabbit mAb (HRP Conjugate) (#9099) Datasheet Without Images Cell Signaling Technol...

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.