

#9182 Store at -20C

Elk-1 Antibody



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H M R Z	Sensitivity: Endogenous	MW (kDa): 47	Source: Rabbit	UniProt ID: #P19419	Entrez-Gene Id: 2002
----------------------------	-------------------------------	-----------------------------------	------------------------	--------------------------	-------------------------------	--------------------------------

Product Usage Information	Application Western Blotting	Dilution 1:1000
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	Elk-1 Antibody detects endogenous levels of total Elk-1 protein. It may cross-react with other Elk/ETS-domain family members.	
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the sequence of human Elk-1. Antibodies are purified by protein A and peptide affinity chromatography.	
Background	The transcription factor Elk-1 is a component of the ternary complex that binds the serum response element (SRE) and mediates gene activity in response to serum and growth factors (1-3). Elk-1 is phosphorylated by MAP kinase pathways at a cluster of S/T motifs at its carboxy terminus; phosphorylation at these sites, particularly Ser383, is critical for transcriptional activation by Elk-1. Elk-1 appears to be a direct target of activated MAP kinase: (a) biochemical studies indicate that Elk-1 is a good substrate for MAP kinase; (b) the kinetics of Elk-1 phosphorylation and activation correlate with MAP kinase activity; (c) interfering mutants of MAP kinase block Elk-1 activation in vivo. Other studies have shown that Elk-1 (Ser383) is also a target of the stress-activated kinase SAPK/JNK (4,5).	
Background References	<ol style="list-style-type: none"> 1. Marais, R. et al. (1993) <i>Cell</i> 73, 381-393. 2. Kortenjann, M. et al. (1994) <i>Mol. Cell. Biol.</i> 14, 4815-4824. 3. Hill, C.S. and Treisman, R. (1995) <i>Cell</i> 80, 199-211. 4. Cavigelli, M. et al. (1995) <i>EMBO J.</i> 14, 5957-5964. 5. Whitmarsh, A.J. et al. (1995) <i>Science</i> 269, 403-407. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or</p>

therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.