

#9441 Store at -20°C

Acetylated-Lysine Antibody


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA
For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB, IP, IHC-P, IF-IC, ChIP, E-P	Reactivity: All	Sensitivity: Endogenous	Source: Rabbit
--	---------------------------	-----------------------------------	--------------------------

Product Usage Information

For optimal ChIP results, use 10 µl of antibody and 10 µg of chromatin (approximately 4 x 10⁶ cells) per IP. This antibody has been validated using SimpleChIP® Enzymatic Chromatin IP Kits.

Application	Dilution
Western Blotting	1:1000
Immunoprecipitation	1:100
Immunohistochemistry (Paraffin)	1:400 - 1:1600
Immunofluorescence (Immunocytochemistry)	1:400 - 1:1600
Chromatin IP	1:50
Peptide ELISA (DELFI)	1:2000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Acetylated-Lysine Antibody detects proteins posttranslationally modified by acetylation on the epsilon-amine groups of lysine residues. The antibody recognizes acetylated lysine in a wide range of sequence contexts. It has been demonstrated to recognize acetylated histones, p53, CBP, PCAF and chemically acetylated BSA. The antibody has been shown to react with as little as 0.04 ng of chemically acetylated BSA while not recognizing up to 25 µg of nonacetylated BSA.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic acetylated lysine-containing peptide. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Acetylation of lysine, like phosphorylation of serine, threonine or tyrosine, is an important reversible modification controlling protein activity. The conserved amino-terminal domains of the four core histones (H2A, H2B, H3, and H4) contain lysines that are acetylated by histone acetyltransferases (HATs) and deacetylated by histone deacetylases (HDACs) (1). Signaling resulting in acetylation/deacetylation of histones, transcription factors, and other proteins affects a diverse array of cellular processes including chromatin structure and gene activity, cell growth, differentiation, and apoptosis (2-6). Recent proteomic surveys suggest that acetylation of lysine residues may be a widespread and important form of post-translational protein modification that affects thousands of proteins involved in control of cell cycle and metabolism, longevity, actin polymerization, and nuclear transport (7,8). The regulation of protein acetylation status is impaired in cancer and polyglutamine diseases (9), and HDACs have become promising targets for anti-cancer drugs currently in development (10).

Background References

- Hassig, C.A. and Schreiber, S.L. (1997) *Curr Opin Chem Biol* 1, 300-8.
- Allfrey, V.G. et al. (1964) *Proc Natl Acad Sci USA* 51, 786-94.
- Liu, L. et al. (1999) *Mol Cell Biol* 19, 1202-9.
- Boyes, J. et al. (1998) *Nature* 396, 594-8.
- Polevoda, B. and Sherman, F. (2002) *Genome Biol* 3, reviews 0006.
- Yoshida, M. et al. (2003) *Prog Cell Cycle Res* 5, 269-78.
- Kim, S.C. et al. (2006) *Mol Cell* 23, 607-18.
- Choudhary, C. et al. (2009) *Science* 325, 834-40.
- Hughes, R.E. (2002) *Curr Biol* 12, R141-3.
- Vigushin, D.M. and Coombes, R.C. (2004) *Curr Cancer Drug Targets* 4, 205-18.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation **IHC-P:** Immunohistochemistry (Paraffin)
IF-IC: Immunofluorescence (Immunocytochemistry) **ChIP:** Chromatin IP **E-P:** Peptide ELISA (DELFI A)

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
 Alexa Fluor is a registered trademark of Life Technologies Corporation.
 All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.