

#9527 Store at -20C

Phospho-cdc25C (Thr48) Antibody


Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 75	Source: Rabbit	UniProt ID: #P30307	Entrez-Gene Id: 995
----------------------------	-------------------------	-----------------------------------	------------------------	--------------------------	-------------------------------	-------------------------------

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Phospho-cdc25C (Thr48) Antibody detects endogenous levels of cdc25C only when phosphorylated at Thr48.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Thr48 of human cdc25C. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Cdc25 is a protein phosphatase responsible for dephosphorylating and activating cdc2, a crucial step in regulating the entry of all eukaryotic cells into mitosis (1). cdc25C is constitutively phosphorylated at Ser216 throughout interphase by c-TAK1, while phosphorylation at this site is DNA damage-dependent at the G2/M checkpoint (2). When phosphorylated at Ser216, cdc25C binds to members of the 14-3-3 family of proteins, sequestering cdc25C in the cytoplasm and thereby preventing premature mitosis (3). The checkpoint kinases Chk1 and Chk2 phosphorylate cdc25C at Ser216 in response to DNA damage (4,5).

Full activation of cdc25C involves phosphorylation at more than 12 different sites by cdc2/cyclin B and Polo-like kinase, and the activity of Pin1, a peptidyl-prolyl isomerase (PPI) (6,7). Pin1 contains a WW domain that binds phospho-Ser/Thr-Pro sites and a catalytic PPI region that induces a cis/trans isomerization on phospho-Ser/Thr-Pro bonds (8). Thr48 and Thr67 of cdc25C interact directly with the WW domain of Pin1 when these sites are phosphorylated (9). Thr48 phosphorylation also mediates binding to CKS/p13SUC1 (10).

Background References

1. Jessus, C. and Ozon, R. (1995) *Prog. Cell Cycle Res.* 1, 215-228.
2. Peng, C.Y. et al. (1997) *Science* 277, 1501-1505.
3. Kumagai, A. and Dunphy, W.G. (1999) *Genes Dev.* 13, 1067-1072.
4. Blasina, A. et al. (1999) *Curr. Biol.* 9, 1-10.
5. Furnari, B. et al. (1999) *Mol. Biol. Cell* 10, 833-845.
6. Izumi, T. and Maller, J.L. (1993) *Mol. Biol. Cell* 4, 1337-1350.
7. Stukenberg, P. T. et al. (2001) *Mol. Cell* 7, 1071-1083.
8. Yaffe, M. B. et al. (1997) *Science* 278, 1957-1960.
9. Lu, P. J. et al. (1999) *Science* 283, 1325-1328.
10. Landrieu, I. et al. (2001) *J. Biol. Chem* 276, 1434-1438.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.