

#9561 Store at -20°C

Phospho- β -Catenin (Ser33/37/Thr41) Antibody



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H M R Mk	Endogenous	92	Rabbit	#P35222	1499

Product Usage Information	Application Western Blotting	Dilution 1:1000
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.	
Specificity / Sensitivity	Phospho- β -Catenin (Ser33/37/Thr41) Antibody detects endogenous levels of β -catenin only when phosphorylated at serines 33, 37 or threonine 41. It does not recognize beta-catenin phosphorylated at other sites.	
Species predicted to react based on 100% sequence homology:	Chicken, Xenopus, Zebrafish, Bovine, Dog, Pig	
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser33, Ser37 and Thr41 of human β -catenin. Antibodies are purified by protein A and peptide affinity chromatography.	
Background	β -catenin is a key downstream effector in the Wnt signaling pathway (1). It is implicated in two major biological processes in vertebrates: early embryonic development (2) and tumorigenesis (3). CK1 phosphorylates β -catenin at Ser45. This phosphorylation event primes β -catenin for subsequent phosphorylation by GSK-3 β (4-6). GSK-3 β destabilizes β -catenin by phosphorylating it at Ser33, Ser37, and Thr41 (7). Mutations at these sites result in the stabilization of β -catenin protein levels and have been found in many tumor cell lines (8).	
Background References	<ol style="list-style-type: none"> 1. Cadigan, K.M. and Nusse, R. (1997) <i>Genes Dev</i> 11, 3286-3305. 2. Wodarz, A. and Nusse, R. (1998) <i>Annu Rev Cell Dev Biol</i> 14, 59-88. 3. Polakis, P. (1999) <i>Curr Opin Genet Dev</i> 9, 15-21. 4. Amit, S. et al. (2002) <i>Genes Dev</i> 16, 1066-76. 5. Liu, C. et al. (2002) <i>Cell</i> 108, 837-47. 6. Yanagawa, S. et al. (2002) <i>EMBO J</i> 21, 1733-42. 7. Yost, C. et al. (1996) <i>Genes Dev</i> 10, 1443-54. 8. Morin, P.J. et al. (1997) <i>Science</i> 275, 1787-90. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.