

#9690
 Store at -20°C

Rab35 Antibody


Cell Signaling
 TECHNOLOGY®

Orders: 877-616-CELL (2355)
 orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
 cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H M R	Endogenous	24	Rabbit	#Q15286	11021

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Rab35 Antibody detects endogeneous levels of total Rab35 protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asp124 of human Rab35 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Rab35 belongs to the small GTPase superfamily. By interacting with its specific GEF or GAP, Rab35 regulates cargo-specific endocytosis at the vesicle recycling step or mediates exocytosis at the exosome docking/tethering step (1-4). During cytokinesis, Rab35 interacts with OCRL phosphatase and is essential for maintaining intercellular bridge stability and abscission by controlling the concentration of phosphatidylinositol 4,5-bisphosphate (PIP2) and SEPT2 localization at the intercellular bridge (5,6). Rab35 also plays a role in actin assembly and the recruitment of Cdc42 and Rac1 to the site of filopodium by its direct interaction with actin-bundling protein fascin and actin-binding protein connecdenn 3 (7-9).

Background References

1. Patino-Lopez, G. et al. (2008) *J Biol Chem* 283, 18323-30.
2. Sato, M. et al. (2008) *EMBO J* 27, 1183-96.
3. Marat, A.L. and McPherson, P.S. (2010) *J Biol Chem* 285, 10627-37.
4. Hsu, C. et al. (2010) *J Cell Biol* 189, 223-32.
5. Kouranti, I. et al. (2006) *Curr Biol* 16, 1719-25.
6. Dambournet, D. et al. (2011) *Nat Cell Biol* 13, 981-8.
7. Zhang, J. et al. (2009) *Science* 325, 1250-4.
8. Marat, A.L. et al. (2012) *Mol Biol Cell* 23, 163-75.
9. Shim, J. et al. (2010) *Mol Cell Biol* 30, 1421-33.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
 All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.