

#9691 Store at -20°C

Nitro-Tyrosine Antibody



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: WB	Reactivity: All	Sensitivity: Endogenous	Source: Rabbit
----------------------------	---------------------------	-----------------------------------	--------------------------

Product Usage Information

To obtain optimal results with this antibody please use PVDF instead of nitrocellulose membranes.

Application	Dilution
Western Blotting	1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Nitro-Tyrosine Antibody detects proteins and peptides containing nitro-tyrosine in a manner independent of the surrounding amino acid sequence. It is a valuable tool for identifying new nitrated proteins as well as for assaying protein nitration and measuring levels of nitrated proteins in tissues and samples. The antibody does not cross-react with unmodified tyrosine or with phospho-tyrosine. (U.S. Patent No's.: 6,441,140; 6,982,318; 7,259,022; 7,344,714; U.S.S.N. 11,484,485; and all foreign equivalents.)

Source / Purification

Polyclonal antibodies are produced by immunizing animals with synthetic nitro-tyrosine-containing peptides. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Nitric oxide (NO) is implicated in carcinogenesis (1), chronic infection, inflammation (2), and neurodegeneration (3). High levels of both superoxide and NO in tissues interact to form peroxynitrite, a potent oxidant that can modify Tyr residues in proteins to form 3-nitro-tyrosine (4). Tyrosine nitration of mitochondrial manganese superoxide dismutase results in loss of enzymatic activity (4). The nitration of p53 at Tyr residues abolishes its capacity for binding to its DNA consensus sequence (5).

Background References

1. Bentz, B.G. et al. (2000) *Head Neck* 22, 64-70.
2. Jaiswal, M. et al. (2000) *Cancer Res* 60, 184-90.
3. Olivenza, R. et al. (2000) *J Neurochem* 74, 785-91.
4. MacMillan-Crow, L.A. et al. (1996) *Proc Natl Acad Sci U S A* 93, 11853-8.
5. Chazotte-Aubert, L. et al. (2000) *Biochem Biophys Res Commun* 267, 609-13.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse **GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner

that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.