

#9931 Store at -20°C

Phospho-Chk1/2 Antibody Sampler Kit

1 Kit (9 x 20 microliters)



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
Phospho-Chk1 (Ser317) (D12H3) XP® Rabbit mAb	12302	20 µl	56 kDa	Rabbit IgG
Phospho-Chk1 (Ser345) (133D3) Rabbit mAb	2348	20 µl	56 kDa	Rabbit IgG
Phospho-Chk1 (Ser296) Antibody	2349	20 µl	56 kDa	Rabbit
Chk1 (2G1D5) Mouse mAb	2360	20 µl	56 kDa	Mouse IgG1
Chk2 (D9C6) Rabbit mAb	6334	20 µl	62 kDa	Rabbit IgG
Phospho-Chk2 (Ser19) Antibody	2666	20 µl	62 kDa	Rabbit
Phospho-Chk2 (Ser33/35) Antibody	2665	20 µl	62 kDa	Rabbit
Phospho-Chk2 (Thr68) (C13C1) Rabbit mAb	2197	20 µl	62 kDa	Rabbit IgG
Phospho-Chk2 (Ser516) Antibody	2669	20 µl	62 kDa	Rabbit
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

The Phospho-Chk1/2 Antibody Sampler Kit offers an economical means to evaluate the phosphorylation status of Chk1 and Chk2 on multiple residues. The kit contains enough primary and secondary antibodies to perform two Western blot experiments with each primary antibody.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Background

Chk1 kinase acts downstream of ATM/ATR kinase and plays an important role in DNA damage checkpoint control, embryonic development, and tumor suppression (1). Activation of Chk1 involves phosphorylation at Ser317 and Ser345 by ATM/ATR, followed by autophosphorylation of Ser296. Activation occurs in response to blocked DNA replication and certain forms of genotoxic stress (2). While phosphorylation at Ser345 serves to localize Chk1 to the nucleus following checkpoint activation (3), phosphorylation at Ser317 along with site-specific phosphorylation of PTEN allows for re-entry into the cell cycle following stalled DNA replication (4). Chk1 exerts its checkpoint mechanism on the cell cycle, in part, by regulating the cdc25 family of phosphatases. Chk1 phosphorylation of cdc25A targets it for proteolysis and inhibits its activity through 14-3-3 binding (5). Activated Chk1 can inactivate cdc25C via phosphorylation at Ser216, blocking the activation of cdc2 and transition into mitosis (6). Centrosomal Chk1 has been shown to phosphorylate cdc25B and inhibit its activation of CDK1-cyclin B1, thereby abrogating mitotic spindle formation and chromatin condensation (7). Furthermore, Chk1 plays a role in spindle checkpoint function through regulation of aurora B and BubR1 (8). Research studies have implicated Chk1 as a drug target for cancer therapy as its inhibition leads to cell death in many cancer cell lines (9).

Chk2 is the mammalian homologue of the budding yeast Rad53 and fission yeast Cds1 checkpoint kinases (5-7). The amino-terminal domain of Chk2 contains a series of seven serine or threonine residues (Ser19, Thr26, Ser28, Ser33, Ser35, Ser50 and Thr68) followed by glutamine (SQ or TQ motif). These are known to be preferred sites for phosphorylation by ATM/ATR kinases (8). Indeed, after DNA damage by ionizing radiation (IR), UV irradiation and DNA replication blocked by hydroxyurea, Thr68 and other sites in this region become phosphorylated by ATM/ATR (9-11). The SQ/TQ cluster domain, therefore, seems to have a regulatory function. Phosphorylation at Thr68 is a prerequisite for the subsequent activation step, which is attributable to autophosphorylation of Chk2 on residues Thr383 and Thr387 in the activation loop of the kinase domain (12).

Background References

1. Liu, Q. et al. (2000) *Genes Dev* 14, 1448-59.
2. Zhao, H. and Piwnicka-Worms, H. (2001) *Mol Cell Biol* 21, 4129-39.
3. Jiang, K. et al. (2003) *J Biol Chem* 278, 25207-17.
4. Martin, S.A. and Ouchi, T. (2008) *Mol Cancer Ther* 7, 2509-16.
5. Chen, M.S. et al. (2003) *Mol Cell Biol* 23, 7488-97.
6. Zeng, Y. et al. (1998) *Nature* 395, 507-10.

7. Löffler, H. et al. (2006) *Cell Cycle* 5, 2543-7.
8. Zachos, G. et al. (2007) *Dev Cell* 12, 247-60.
9. Garber, K. (2005) *J Natl Cancer Inst* 97, 1026-8.
10. Allen, J.B. et al. (1994) *Genes Dev* 8, 2401-15.
11. Weinert, T.A. et al. (1994) *Genes Dev* 8, 652-65.
12. Murakami, H. and Okayama, H. (1995) *Nature* 374, 817-9.
13. Kastan, M.B. and Lim, D.S. (2000) *Nat Rev Mol Cell Biol* 1, 179-86.
14. Matsuoka, S. et al. (2000) *Proc Natl Acad Sci U S A* 97, 10389-94.
15. Melchionna, R. et al. (2000) *Nat Cell Biol* 2, 762-5.
16. Ahn, J.Y. et al. (2000) *Cancer Res* 60, 5934-6.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.