

#11828 Store at -20C

Arrestin 1/S-Arrestin Antibody

**Cell Signaling**
TECHNOLOGY®**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H M R	Endogenous	52	Rabbit	#P10523	6295

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Arrestin 1/S-Arrestin Antibody recognizes endogenous levels of total Arrestin 1/S-Arrestin protein.

Species predicted to react based on 100% sequence homology:

Monkey

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala163 of human Arrestin 1/S-Arrestin protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Arrestin proteins function as negative regulators of G protein-coupled receptor (GPCR) signaling. Cognate ligand binding stimulates GPCR phosphorylation, which is followed by binding of arrestin to the phosphorylated GPCR and the eventual internalization of the receptor and desensitization of GPCR signaling (1). Four distinct mammalian arrestin proteins are known. Arrestin 1 (also known as S-arrestin) and arrestin 4 (X-arrestin) are localized to retinal rods and cones, respectively. Arrestin 2 (also known as β-arrestin 1) and arrestin 3 (β-arrestin 2) are ubiquitously expressed and bind to most GPCRs (2). β-arrestins function as adaptor and scaffold proteins and play important roles in other processes, such as recruiting c-Src family proteins to GPCRs in Erk activation pathways (3,4). β-arrestins are also involved in some receptor tyrosine kinase signaling pathways (5-8). Additional evidence suggests that β-arrestins translocate to the nucleus and help regulate transcription by binding transcriptional cofactors (9,10). Arrestin 1/S-Arrestin inactivates rhodopsin-mediated signaling through binding to the light-activated phosphorylated rod photoreceptor, thereby preventing coupling to transducin, rhodopsin's cognate G protein (11). Research studies have proposed that mutations in the Arrestin 1/S-Arrestin gene are linked to Oguchi disease (12,13) and Retinitis Pigmentosa (14).

Background References

- Shenoy, S.K. and Lefkowitz, R.J. (2005) *Sci STKE* 2005, cm10.
- Lefkowitz, R.J. and Shenoy, S.K. (2005) *Science* 308, 512-7.
- Luttrell, L.M. et al. (1999) *Science* 283, 655-61.
- Luttrell, L.M. et al. (1999) *Curr Opin Cell Biol* 11, 177-83.
- Luttrell, L.M. and Lefkowitz, R.J. (2002) *J Cell Sci* 115, 455-65.
- Waters, C. et al. (2004) *Semin Cell Dev Biol* 15, 309-23.
- Lefkowitz, R.J. and Whalen, E.J. (2004) *Curr Opin Cell Biol* 16, 162-8.
- Waters, C.M. et al. (2005) *Cell Signal* 17, 263-77.
- Kang, J. et al. (2005) *Cell* 123, 833-47.
- Ma, L. and Pei, G. (2007) *J Cell Sci* 120, 213-8.
- Gurevich, V.V. et al. (2011) *Prog Retin Eye Res* 30, 405-30.
- Yamada, T. et al. (1999) *Ophthalmic Genet* 20, 117-20.
- Waheed, N.K. et al. (2012) *Mol Vis* 18, 1253-9.
- Nakazawa, M. et al. (1998) *Arch Ophthalmol* 116, 498-501.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.