

#11831 Store at -20C

## WBP2 Antibody

Cell Signaling  
TECHNOLOGY®Orders: 877-616-CELL (2355)  
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H M R Mk	Endogenous	32	Rabbit	#Q969T9	23558

## Product Usage Information

## Application

Western Blotting

## Dilution

1:1000

## Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

## Specificity / Sensitivity

WBP2 Antibody recognizes endogenous levels of total WBP2 protein.

## Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asn239 of human WBP2 protein. Antibodies are purified by protein A and peptide affinity chromatography.

## Background

WW-domain binding protein 2 (WBP2) is an adaptor protein first identified in a screen for proteins that interact with YAP1 (1). WBP2 was subsequently shown to bind other WW domain-containing proteins, including TAZ and NEDD4-like ubiquitin-protein ligases (2-5). There is strong evidence for a conserved functional role for WBP2 in the Hippo kinase tumor suppressor pathway. WBP2 interaction is required for the oncogenic properties of TAZ in breast cancer cells (4) and the growth-promoting properties of the YAP homolog Yorkie in *Drosophila* (6). *In vitro* studies have also suggested that WBP2 may act as a transcriptional co-activator of both estrogen and progesterone receptors (7); this interaction was shown to be dependent upon tyrosine phosphorylation of WBP2 by c-Src and c-Yes kinases (8).

## Background References

1. Chen, H.I. and Sudol, M. (1995) *Proc Natl Acad Sci U S A* 92, 7819-23.
2. Chen, H.I. et al. (1997) *J Biol Chem* 272, 17070-7.
3. McDonald, C.B. et al. (2011) *Biochemistry* 50, 9616-27.
4. Chan, S.W. et al. (2011) *Oncogene* 30, 600-10.
5. Jolliffe, C.N. et al. (2000) *Biochem J* 351 Pt 3, 557-65.
6. Zhang, X. et al. (2011) *Cell Death Differ* 18, 1346-55.
7. Dhananjayan, S.C. et al. (2006) *Mol Endocrinol* 20, 2343-54.
8. Lim, S.K. et al. (2011) *FASEB J* 25, 3004-18.

## Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

## Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

## Applications Key

WB: Western Blotting

## Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster  
X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse  
GP: Guinea Pig Rab: rabbit All: all species expected

## Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.