

#11867 Store at -20C

NBC1/SLC4A4 Antibody

**Cell Signaling**
TECHNOLOGY®**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M Mk	Endogenous	135-145	Rabbit	#Q9Y6R1	8671

Product Usage Information**Application**Western Blotting
Immunoprecipitation**Dilution**1:1000
1:50**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

NBC1/SLC4A4 Antibody recognizes endogenous levels of total NBC1 protein. This antibody also cross-reacts with 42 and 62 kDa proteins of unknown origin in some cell lines.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human NBC1 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

NBC1/SLC4A4 is an electrogenic sodium bicarbonate cotransporter. It mediates the coupled movement of Na⁺ and HCO₃⁻ ions across the basolateral membrane of cells. NBC1 regulates secretion and absorption of bicarbonate and modulates intracellular pH (1-3). There are five isoforms of NBC1 produced by alternative splicing including the most studied isoforms, 1 and 2, which have different N termini. The N terminus of isoform 1 contains multiple consensus phosphorylation sites for various kinases, such as PKA, PKC, and CK II, and may play a regulatory role in the biological function of NBC1 (4). Interaction with carbonic anhydrase II, IV, and IX regulates the transporter activity of NBC1 (5-7). NBC1 is expressed in various tissues with the exception of isoform 2, which is mainly expressed in kidney proximal tubules (1). Isoform 1 is expressed in the pancreas and corneal endothelium, and at low levels in other tissues including heart, skeletal muscle, liver, and prostate (4). Research studies have shown that mutations in the NBC1 gene are linked to proximal renal tubular acidosis with ocular abnormalities (also known as renal tubular acidosis II).

Background References

1. Burnham, C.E. et al. (1997) *J Biol Chem* 272, 19111-4.
2. Amlal, H. et al. (1998) *J Biol Chem* 273, 16810-5.
3. Choi, I. et al. (1999) *Am J Physiol* 276, C576-84.
4. Abuladze, N. et al. (1998) *J Biol Chem* 273, 17689-95.
5. Alvarez, B.V. et al. (2003) *Biochemistry* 42, 12321-9.
6. Loisel, F.B. et al. (2004) *Am J Physiol Cell Physiol* 286, C1423-33.
7. Svastova, E. et al. (2012) *J Biol Chem* 287, 3392-402.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key**WB:** Western Blotting **IP:** Immunoprecipitation**Cross-Reactivity Key**

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
 All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.