

#11878 Store at -20C

UBE2S (D5H9H) Rabbit mAb**Cell Signaling**
TECHNOLOGY®**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R Mk	Endogenous	26	Rabbit IgG	#Q16763	27338

Product Usage Information**Application****Dilution**

Western Blotting

1:1000

Immunoprecipitation

1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

UBE2S (D5H9H) Rabbit mAb recognizes endogenous levels of total UBE2S protein.

Species predicted to react based on 100% sequence homology:

Bovine, Dog

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human UBE2S protein.

Background

Protein ubiquitination requires the concerted action of the E1, E2, and E3 ubiquitin-conjugating enzymes. Ubiquitin is first activated through ATP-dependent formation of a thiol ester with ubiquitin-activating enzyme E1. The activated ubiquitin is then transferred to a thiol group of ubiquitin-carrier enzyme E2. The final step is the transfer of ubiquitin from E2 to an ε-amino group of the target protein lysine residue, which is mediated by ubiquitin-ligase enzyme E3 (1).

The human anaphase promoting complex (APC/C) is a large macromolecular E3 ligase complex that is largely responsible for timely progression through mitosis via the sequential targeting of cell cycle regulators for proteasomal degradation. Recent work has revealed that APC/C substrates are marked for proteasomal degradation during cell cycle progression through the covalent assembly of Lys11-linked ubiquitin chains, which occurs through a priming phase and an elongation phase (2-5). The APC/C utilizes, in part, the UBE2C/UBCH10 E2 enzyme to prime substrates for degradation through the covalent attachment of short Lys11-linked chains (3,6). The Lys11-specific elongating E2 enzyme, UBE2S/E2-EPF, extends these short chains into long Lys11-linked ubiquitin chains on APC/C bound substrates (2,3,7). In addition to the well-established biochemical role for UBE2S in cell cycle regulation, researchers have found evidence that this enzyme is overexpressed in many types of human cancer (8), and has been implicated in hypoxia signaling (9,10). Indeed, UBE2S has been reported by researchers to associate with VHL and to target it for proteasomal degradation, thereby stabilizing HIF-1α (9).

Background References

1. Hershko, A. (1988) *J Biol Chem* 263, 15237-40.
2. Williamson, A. et al. (2009) *Proc Natl Acad Sci U S A* 106, 18213-8.
3. Jin, L. et al. (2008) *Cell* 133, 653-65.
4. Wu, T. et al. (2010) *Proc Natl Acad Sci U S A* 107, 1355-60.
5. Song, L. and Rape, M. (2010) *Mol Cell* 38, 369-82.
6. Summers, M.K. et al. (2008) *Mol Cell* 31, 544-56.
7. Wickliffe, K.E. et al. (2011) *Cell* 144, 769-81.
8. Tedesco, D. et al. (2007) *Neoplasia* 9, 601-13.
9. Jung, C.R. et al. (2006) *Nat Med* 12, 809-16.
10. Roos, F.C. et al. (2011) *Am J Pathol* 178, 853-60.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.