

#11990 Store at -20°C

Tensin 2 Antibody

Cell Signaling
TECHNOLOGY®Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R	Endogenous	145-155	Rabbit	#Q63HR2	23371

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Immunoprecipitation

1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Tensin 2 Antibody recognizes endogenous levels of total Tensin 2 protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Arg155 of human Tensin 2 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Tensin 2 belongs to the Tensin family of cytoskeletal proteins that includes Tensin 1-3 and Cten, which couple integrins to the actin cytoskeleton (1). Tensin proteins contain SH2 and phosphotyrosine binding (PTB) domains, which enable interaction with diverse signaling molecules and proteins. Tensin family proteins play important roles in signal transduction, cell proliferation, and motility (2-5).

Tensin 2 is localized to focal adhesions of various tissues with highest expression in the heart, kidney, and liver (6,7). Tensin 2 inhibits Akt/PKB signaling via a phosphatase tensin-type domain (8). However, Tensin 2 also mediates thrombopoietin/c-Mpl signaling, which promotes Akt signaling (9). Interaction with Tensin 2 is essential for the tumor suppressor function of Deleted in Cancer 1 (DLC1) (6,10,11).

Background References

- Lo, S.H. et al. (1994) *Bioessays* 16, 817-23.
- Lo, S.H. et al. (1994) *J Cell Biol* 125, 1067-75.
- Chen, H. and Lo, S.H. (2003) *Biochem J* 370, 1039-45.
- Katz, M. et al. (2007) *Nat Cell Biol* 9, 961-9.
- Chen, H. et al. (2002) *Proc Natl Acad Sci U S A* 99, 733-8.
- Clark, K. et al. (2010) *J Cell Biochem* 109, 808-17.
- Hafizi, S. et al. (2002) *Biochem Biophys Res Commun* 299, 793-800.
- Hafizi, S. et al. (2005) *FASEB J* 19, 971-3.
- Jung, A.S. et al. (2011) *Cell Cycle* 10, 1838-44.
- Chan, L.K. et al. (2009) *PLoS One* 4, e5572.
- Chen, L. et al. (2012) *J Biol Chem* 287, 26104-14.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
 All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.