

#12117 Store at -20°C

ITCH (D8Q6D) Rabbit mAb



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H M R	Endogenous	105	Rabbit IgG	#Q96J02	83737

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:200

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

ITCH (D8Q6D) Rabbit mAb recognizes endogenous levels of total ITCH protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asp125 of human ITCH protein.

Background

ITCH is a HECT domain-containing E3 ubiquitin ligase, first identified in genetic studies of the mouse *agouti* locus, in which mutations result in characteristic coat color changes. One particular *agouti* mutation (non-*agouti*-lethal 18H) is notable for the development of immunological defects not observed in other *agouti* mutant mice; these include lymphoid hyperplasia and chronic stomach, lung and skin inflammation (manifest as constant itching). The 18H *agouti* mutation was traced to a chromosomal inversion that disrupted expression of an adjacent gene in the *agouti* locus, subsequently termed *Itch* to reflect the chronic itching phenotype (1-3). Further characterizations revealed that *Itch* encoded a NEDD4-like E3-ubiquitin ligase capable of catalyzing Lys29, Lys48, and/or Lys63-linked ubiquitination of target proteins, leading to their degradation by the proteasome pathway (4-6). The distinct phenotypes of *Itch* mutant mice led to the identification of an important regulatory role for ITCH-mediated ubiquitination in inflammatory signaling pathways. For example, ITCH-mediated ubiquitination of the transcription factor JunB was shown to play a direct inhibitory role in regulating expression of the proinflammatory cytokine IL-4. ITCH-null T lymphocytes consequently exhibit increased production of IL-4, leading to biased differentiation of naive CD4⁺ cells towards the proinflammatory Th2 lineage (7). In accordance with the findings from mutant *Itch* mouse models, a genetic linkage study in humans identified loss-of-function mutations in *ITCH* as a direct cause of syndromic multisystem autoimmune disease (SMAD) (8). Notably, targets of ITCH-mediated ubiquitination are not restricted to immune signaling pathways. For example, key mediators of the Hedgehog (9,10), Wnt/β-catenin (11), Hippo (12), and Notch signaling pathways (13,14) have been identified as important targets of ITCH-mediated ubiquitination (2).

Background References

1. Matesic, L.E. et al. (2008) *Curr Top Microbiol Immunol* 321, 185-200.
2. Melino, G. et al. (2008) *Cell Death Differ* 15, 1103-12.
3. Perry, W.L. et al. (1998) *Nat Genet* 18, 143-6.
4. Chastagner, P. et al. (2006) *EMBO Rep* 7, 1147-53.
5. Lee, T.L. et al. (2008) *Biochem Biophys Res Commun* 375, 326-30.
6. Ahmed, N. et al. (2011) *Nat Immunol* 12, 1176-83.
7. Fang, D. et al. (2002) *Nat Immunol* 3, 281-7.
8. Lohr, N.J. et al. (2010) *Am J Hum Genet* 86, 447-53.
9. Di Marcotullio, L. et al. (2006) *Nat Cell Biol* 8, 1415-23.
10. Di Marcotullio, L. et al. (2011) *Oncogene* 30, 65-76.
11. Wei, W. et al. (2012) *Mol Cell Biol* 32, 3903-12.
12. Salah, Z. et al. (2011) *Cancer Res* 71, 2010-20.
13. Qiu, L. et al. (2000) *J Biol Chem* 275, 35734-7.
14. McGill, M.A. and McGlade, C.J. (2003) *J Biol Chem* 278, 23196-203.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.