

#12283 Store at -20C

CARD9 Antibody

Cell Signaling
TECHNOLOGY®Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB, IP	M	Endogenous	64	Rabbit	#Q9H257	64170

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Immunoprecipitation

1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

CARD9 Antibody recognizes endogenous levels of total CARD9 protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding His478 of mouse CARD9 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

CARD9 is a caspase recruitment domain (CARD)-containing adaptor protein expressed by myeloid cells (1,2). It is required for antifungal immunity downstream of pathogen detection by C-type lectin receptors (CLRs) such as Dectin-1 (3,4). Recognition of carbohydrates on fungal cell walls by CLRs leads to activation of the tyrosine kinase Syk, followed by activation of PKCδ (5,6). PKCδ phosphorylates CARD9, enabling the assembly of a complex containing CARD9 and Bcl10 (6). This complex activates NF-κB, resulting in upregulation of inflammatory cytokines important for initiation of adaptive immunity (3,4,6,7). CARD9 was also shown to be important for the induction of IL-1β, downstream of the viral nucleic acid sensor RIG-I, as well as for the generation of reactive oxygen species important for bacterial killing by macrophages (2,8).

Background References

1. Bertin, J. et al. (2000) *J Biol Chem* 275, 41082-6.
2. Hsu, Y.M. et al. (2007) *Nat Immunol* 8, 198-205.
3. Gross, O. et al. (2006) *Nature* 442, 651-6.
4. Bi, L. et al. (2010) *J Biol Chem* 285, 25969-77.
5. Rogers, N.C. et al. (2005) *Immunity* 22, 507-17.
6. Strasser, D. et al. (2012) *Immunity* 36, 32-42.
7. LeibundGut-Landmann, S. et al. (2007) *Nat Immunol* 8, 630-8.
8. Wu, W. et al. (2009) *Nat Immunol* 10, 1208-14.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key

H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster
X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse
GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.