e at -20C	Pumilio 1 Antibody		Cell Signaling TECHNOLOGY®	
Store		Orders:	877-616-CELL (2355) orders@cellsignal.com	
322		Support:	877-678-TECH (8324)	
1232		Web:	info@cellsignal.com cellsignal.com	
#		3 Trask Lane Danvers Ma	ssachusetts 01923 USA	

For Research Use Only. Not for Use in Diagnostic Procedures.

	tivity: Sensitivity: M Endogenous	MW (kDa): 140	Source: Rabbit	UniProt ID: #Q14671	Entrez-Gene Id: 9698			
Product Usage Information	Application Western Blotting							
	Immunoprecipitation			1:100				
Storage	Supplied in 10 mM sodi	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.						
Specificity / Sensitivity	Pumilio 1 Antibody reco	Pumilio 1 Antibody recognizes endogenous levels of total pumilio 1 protein.						
Species predicted to react based on 100% sequence homology:	Rat, Monkey	Rat, Monkey						
Source / Purification	Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly169 of human pumilio 1 protein. Antibodies are purified by protein A and peptide affinity chromatography.							
Background	conserved RNA binding binding to the 3'-UTR of regulation of genes invo	Pumilio 1 and Pumilio 2 (PUM1, PUM2, or Pumilio homolog 1 and 2, respectively) are evolutionarily conserved RNA binding proteins that are thought to repress translation and stability of mRNA targets by binding to the 3'-UTR of specific RNA sequences (1). Pumilio proteins have been implicated in the regulation of genes involved in embryogenesis and germline cell development (2). Research studies have shown that PUM2 may have a role in neural stem cell fate decisions (3).						
Background References	 Weidmann, C.A. and Goldstrohm, A.C. (2012) Mol Cell Biol 32, 527-40. Spassov, D.S. and Jurecic, R. (2003) IUBMB Life 55, 359-66. Vessey, J.P. et al. (2012) Cell Stem Cell , . 							
Species Reactivity	Species reactivity is dete	ermined by testing i	n at least one appro	ved application (e.g., we	estern blot).			
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.							
Applications Key	WB: Western Blotting IP: Immunoprecipitation							
Cross-Reactivity Key	X: Xenopus Z: zebrafish	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected						
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.						
Limited Uses	following terms apply to I conditions that are in add	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.						

Pumilio 1 Antibody (#12322) Datasheet Without Images Cell Signaling Technology

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.