

#12436 Store at -20C

FIP200 (D10D11) Rabbit mAb**Cell Signaling**
TECHNOLOGY®**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

| Applications: | Reactivity: | Sensitivity: | MW (kDa): | Source/Isotype: | UniProt ID: | Entrez-Gene Id: |
|---------------|-------------|--------------|-----------|-----------------|-------------|-----------------|
| WB, IP | H M | Endogenous | 200 | Rabbit IgG | #Q8TDY2 | 9821 |

Product Usage Information**Application**Western Blotting
Immunoprecipitation**Dilution**1:1000
1:100**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

FIP200 (D10D11) Rabbit mAb recognizes endogenous levels of total FIP200 protein.

Species predicted to react based on 100% sequence homology:

Monkey

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human FIP200 protein.

Background

FIP200 (FAK family kinase-interacting protein of 200 kDa) was identified in a two-hybrid screen with the tyrosine kinase Pyk2 and can inhibit Pyk2 kinase activity as well as related family members (1). FIP200 was later independently identified in a multi-drug resistance screen and named RB1CC1 (RB1-inducible coiled-coil 1) due to its induction by cytotoxic stress and RB1 expression regulation (2). FIP200 function has been linked to apoptosis, cell cycle progression, cell growth, and migration (reviewed in 3). FIP200 has also recently been shown to interact with ULK1 and is required for autophagosome formation (4). FIP200 is part of an ULK1 complex along with Atg13 that is regulated by mTOR and is required for starvation induced autophagy (5-7).

Background References

1. Ueda, H. et al. (2000) *J Cell Biol* 149, 423-30.
2. Chano, T. et al. (2002) *Oncogene* 21, 1295-8.
3. Gan, B. and Guan, J.L. (2008) *Cell Signal* 20, 787-94.
4. Hara, T. et al. (2008) *J Cell Biol* 181, 497-510.
5. Hosokawa, N. et al. (2009) *Mol Biol Cell* 20, 1981-91.
6. Jung, C.H. et al. (2009) *Mol Biol Cell* 20, 1992-2003.
7. Ganley, I.G. et al. (2009) *J Biol Chem* 284, 12297-305.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key**WB:** Western Blotting **IP:** Immunoprecipitation**Cross-Reactivity Key**

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
 All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.