

#12551 Store at -20°C

**mGluR1 (D5H10) Rabbit mAb****Cell Signaling**  
TECHNOLOGY®**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP, IHC-P, IF-F	H M R	Endogenous	145, >300	Rabbit IgG	#Q13255	2911

**Product Usage Information****Application**

Western Blotting

Immunoprecipitation

Immunohistochemistry (Paraffin)

Immunofluorescence (Frozen)

**Dilution**

1:1000

1:50

1:100 - 1:400

1:400 - 1:1600

**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

**Specificity / Sensitivity**

mGluR1 (D5H10) Rabbit mAb recognizes endogenous levels of total mGluR1 protein.

**Source / Purification**

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Leu1105 of human mGluR1 protein.

**Background**

Metabotropic glutamate receptor 1 (mGluR1) is a G protein-coupled receptor (GPCR) for the neurotransmitter glutamate in the mammalian brain. Unlike ionotropic receptors, metabotropic receptors do not form an ion channel pore themselves but are indirectly linked to ion channels (1). Both mGluR1 and mGluR5 are coupled to phospholipase C and activate inositol phospholipid metabolism via G protein-mediated mechanisms. Upon phosphatidylinositol activation, the second messenger calcium is released and generates a calcium-activated chloride current. Metabotropic glutamate receptors other than mGluR1 and mGluR5 inhibit adenylate cyclase (1-3). mGluR1 does not share sequence homology with conventional GPCRs (1). mGluR1 forms a homodimer and is linked to synaptic plasticity, as well as long-term potentiation and long-term depression. Furthermore, mGluR1 is a potential therapeutic target for various psychiatric and neurological diseases, including schizophrenia, epilepsy, and Parkinson and Alzheimer diseases (4-6).

**Background References**

1. Pin, J.P. et al. (1994) *EMBO J* 13, 342-8.
2. Sugiyama, H. et al. (1987) *Nature* 325, 531-3.
3. Hermans, E. and Challiss, R.A. (2001) *Biochem J* 359, 465-84.
4. Niswender, C.M. et al. (2005) *Curr Top Med Chem* 5, 847-57.
5. Pellicciari, R. and Costantino, G. (1999) *Curr Opin Chem Biol* 3, 433-40.
6. Olive, M.F. (2009) *Curr Drug Abuse Rev* 2, 83-989.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key****WB:** Western Blotting **IP:** Immunoprecipitation **IHC-P:** Immunohistochemistry (Paraffin)  
**IF-F:** Immunofluorescence (Frozen)**Cross-Reactivity Key****H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster  
**X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse  
**GP:** Guinea Pig **Rab:** rabbit **All:** all species expected**Trademarks and Patents**Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.  
All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.