

#12660 Store at -20C

CABIN1 (D2B9F) Rabbit mAb**Cell Signaling**
TECHNOLOGY®**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, IP	H	Endogenous	230	Rabbit IgG	#Q9Y6J0	23523

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Immunoprecipitation

1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity / Sensitivity

CABIN1 (D2B9F) Rabbit mAb recognizes endogenous levels of total CABIN1 protein.

Species predicted to react based on 100% sequence homology:

Monkey

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human CABIN1 protein.

Background

Calcineurin binding protein CABIN1 was originally identified as an inhibitor of the calcium-dependent serine/threonine phosphatase, calcineurin. CABIN1 inhibits calcineurin signaling in T cells, regulating T cell receptor (TCR) signaling, transcription, and apoptosis (1-4). CABIN1 represses myocyte enhancer factor 2 (MEF2)-mediated transcription by recruiting chromatin remodeling enzymes (5), and also negatively regulates the activity of the tumor suppressor p53 (6). In response to genotoxic stress, CABIN1 is degraded and releases its inhibition of p53, allowing p53 to elicit cellular stress responses (7). CABIN1 is also involved in regulation of chromatin structure as part of the highly conserved HIRA/UBN1/CABIN1/ASF1A (HUCA) histone chaperone complex (8,9).

Background References

1. Sun, L. et al. (1998) *Immunity* 8, 703-11.
2. Youn, H.D. and Liu, J.O. (2000) *Immunity* 13, 85-94.
3. Liu, W. et al. (2001) *Eur J Immunol* 31, 1757-64.
4. Kim, M.J. et al. (2002) *Proc Natl Acad Sci U S A* 99, 9870-5.
5. Jang, H. et al. (2007) *J Biol Chem* 282, 11172-9.
6. Jang, H. et al. (2009) *Nat Struct Mol Biol* 16, 910-5.
7. Choi, S.Y. et al. (2013) *Nucleic Acids Res* 41, 2180-2190.
8. Rai, T.S. et al. (2011) *Mol Cell Biol* 31, 4107-18.
9. Tang, Y. et al. (2012) *Biochemistry* 51, 2366-77.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key**WB:** Western Blotting **IP:** Immunoprecipitation**Cross-Reactivity Key**

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster
X: Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse
GP: Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.